




New Day in Medicine
Новый День в Медицине

NDM



TIBBIYOTDA YANGI KUN

Ilmiy referativ, marifiy-ma'naviy jurnal



AVICENNA-MED.UZ



ISSN 2181-712X.
EiSSN 2181-2187

3 (53) 2023

Сопредседатели редакционной коллегии:

**Ш. Ж. ТЕШАЕВ,
А. Ш. РЕВИШВИЛИ**

Ред. коллегия:

М.И. АБДУЛЛАЕВ
А.А. АБДУМАЖИДОВ
А.Ш. АБДУМАЖИДОВ
Р.Б. АБДУЛЛАЕВ
М.М. АКБАРОВ
Х.А. АКИЛОВ
М.М. АЛИЕВ
С.Ж. АМИНОВ
Ш.Э. АМОНОВ
Ш.М. АХМЕДОВ
Ю.М. АХМЕДОВ
Т.А. АСКАРОВ
Ж.Б. БЕКНАЗАРОВ (главный редактор)
Е.А. БЕРДИЕВ
Б.Т. БУЗРУКОВ
Р.К. ДАДАБАЕВА
М.Н. ДАМИНОВА
К.А. ДЕХКОНОВ
Э.С. ДЖУМАБАЕВ
А.Ш. ИНОЯТОВ
С. ИНДАМИНОВ
А.И. ИСКАНДАРОВ
С.И. ИСМОИЛОВ
Э.Э. КОБИЛОВ
Д.М. МУСАЕВА
Т.С. МУСАЕВ
Ф.Г. НАЗИРОВ
Н.А. НУРАЛИЕВА
Б.Т. РАХИМОВ
Ш.И. РУЗИЕВ
С.А. РУЗИБОЕВ
С.А. ГАФФОРОВ
Ж.Б. САТТАРОВ
Б.Б. САФОЕВ (отв. редактор)
И.А. САТИВАЛДИЕВА
Д.И. ТУКСАНОВА
М.М. ТАДЖИЕВ
А.Ж. ХАМРАЕВ
А.М. ШАМСИЕВ
А.К. ШАДМАНОВ
Н.Ж. ЭРМАТОВ
Б.Б. ЕРГАШЕВ
Н.Ш. ЕРГАШЕВ
И.Р. ЮЛДАШЕВ
Д.Х. ЮЛДАШЕВА
А.С. ЮСУПОВ
М.Ш. ХАКИМОВ
К.А. ЕГЕЗАРЯН (Россия)
DONG JINCHENG (Китай)
КУЗАКОВ В.Е. (Россия)
Я. МЕЙЕРНИК (Словакия)
В.А. МИТИШ (Россия)
В.И. ПРИМАКОВ (Беларусь)
О.В. ПЕШИКОВ (Россия)
А.А. ПОТАПОВ (Россия)
А.А. ТЕПЛОВ (Россия)
Т.Ш. ШАРМАНОВ (Казахстан)
А.А. ЩЕГОЛОВ (Россия)
Prof. Dr. KURBANHAN MUSLUMOV (Azerbaijan)
Prof. Dr. DENIZ UYAK (Germany)

www.bsmi.uz

<https://newdaymedicine.com>

E: ndmuz@mail.ru

Тел: +99890 8061882

**ТИББИЁТДА ЯНГИ КУН
НОВЫЙ ДЕНЬ В МЕДИЦИНЕ
NEW DAY IN MEDICINE**

Илмий-рефератив, маънавий-маърифий журнал

Научно-реферативный,

духовно-просветительский журнал

УЧРЕДИТЕЛИ:

**БУХАРСКИЙ ГОСУДАРСТВЕННЫЙ
МЕДИЦИНСКИЙ ИНСТИТУТ
ООО «ТИББИЁТДА ЯНГИ КУН»**

Национальный медицинский
исследовательский центр хирургии имени
А.В. Вишневского является генеральным
научно-практическим
консультантом редакции

Журнал был включен в список журнальных
изданий, рецензируемых Высшей
Аттестационной Комиссией
Республики Узбекистан
(Протокол № 201/03 от 30.12.2013 г.)

РЕДАКЦИОННЫЙ СОВЕТ:

М.М. АБДУРАХМАНОВ (Бухара)
Г.Ж. ЖАРЫЛКАСЫНОВА (Бухара)
А.Ш. ИНОЯТОВ (Ташкент)
Г.А. ИХТИЁРОВА (Бухара)
Ш.И. КАРИМОВ (Ташкент)
У.К. КАЮМОВ (Ташкент)
Ш.И. НАВРУЗОВА (Бухара)
А.А. НОСИРОВ (Ташкент)
А.Р. ОБЛОКУЛОВ (Бухара)
Б.Т. ОДИЛОВА (Ташкент)
Ш.Т. УРАКОВ (Бухара)

3 (53)

2023

март

Received: 20.02.2023, Accepted: 25.02.2023, Published: 15.03.2023.

UDK 613.2. 614.31:633

ASSESSMENT OF PHYSICAL DEVELOPMENT FROM A HYGIENIC POINT OF VIEW AND DETERMINE SOME FACTORS AND INFLUENCES ON IT IN CHILDREN FROM 4 TO 7 YEARS OLD IN BUKHARA

Kosimov X.O. <https://orcid.org/0000-0002-2870-3248>

Samadova X.S. <https://orcid.org/0009-0006-6535-8849>

Bukhara State Medical Institute named after Abu Ali Ibn Sina Uzbekistan Bukhara, A.Navoiist. 1

Tel: +998(65) 223-00-50 e-mail: info@bsmi.uz

✓ *Resume*

The purpose of the study is to identify gender, age and constitutional features of physical development and motor qualities in children aged 4- 7 years in Bukhara.

The purpose of the study is to identify gender, age and constitutional features of physical development and motor qualities in children aged 4- 7 years in Bukhara.

Also, an increase even in boys with high and above average levels of physical development is noted at 6 and 7 years old and slightly at 4 years old.

Keywords: physical development, endurance, motor activity, chest.

ОЦЕНКА ФИЗИЧЕСКОГО РАЗВИТИЯ С ГИГИЕНИЧЕСКОЙ ТОЧКИ ЗРЕНИЯ И ОПРЕДЕЛИТЬ НЕКОТОРЫЕ ФАКТОРЫ И ВЛИЯЮЩИЕ НА НЕГОДЕТИ ОТ 4 ДО 7 ЛЕТг. БУХАРЫ

Косимов Х.О., Самадова Х.С.

Бухарский государственный медицинский институт имени Абу Али Ибн Сины, Узбекистан

✓ *Резюме*

Цель исследования является выявить половые возрастные и конституциональные особенности физического развития и двигательных качеств у детей возрасте 4- 7 лет в г Бухары.

Изучение интегральной показателя физического развития у мальчиков 4-1 лет выявило наибольшую их долю со средним уровнем развития во всех возрастах. При этом в 5 лет таких детей становится меньше, чем как у девочек, а с уровнями выше среднего и высоким, напротив, больше (до 39,8%).

Так же увеличение даже мальчиков с уровнями физического развития высоким и выше среднего отмечается в 6 и 7 лет и незначительно - в 4 года.

Ключевые слова: физическое развитие, выносливость, двигательной активность, грудной клетки.

BUXORO SHAHRIDAGI 4 YOSHDAN 7 YOSHGACHA BO‘LGAN BOLALARNING JISMONIY RIVOJLANISHINI GIGIYENIK NUQTAI NAZARDAN BAHOLASH VA UNGA AYRIM OMILLAR VA TA’SIRLARNI ANIQLASH

Qosimov X.O., Samadova X.S.

Abu Ali Ibn Sino nomidagi Buxoro davlat tibbiyot instituti, O‘zbekiston

✓ *Rezyume*

Tadqiqot maqsadining Buxoro shahridagi 4-7 yoshli bolalarning jinsi, yoshi va jismoniy rivojlanishi va harakat sifatlarining konstitutsiyaviy xususiyatlarini aniqlash.4-7 yoshdagi o'g'ilbolalarning jismoniy rivojlanishining integral ko'rsatkichini o'rganish ularning barcha yoshdagi o'rtacha rivojlanish darajasiga ega bo'lgan eng katta qismini aniqladi. Shu bilan birga, 5

yoshda bunday bolalar o'rtacha va yuqori darajadan yuqori bo'lgan qizlarga qaraganda kamroq, aksincha, ko'proq (39,8% gacha).

Shuningdek, jismoniy rivojlanish darajasi yuqori va o'rtachadan yuqori bo'lgan o'g'il bolalarda ham o'sish 6 va 7 yoshda va 4 yoshda biroz kuzatiladi.

Kalit so'zlar: jismoniy rivojlanish, chidamlilik, vosita faolligi, ko'krak qafasi.

Relevance

Currently, at the global level, there has been a steady trend of some deterioration in physical health and a decrease in the functional capabilities of their body [9,10,11] negative trends are also observed in the physical development of children, adolescents and young people, which are manifested by a decrease in body length and weight (desekliration) and an increase in the disharmony of physical development [1,6,9,10].

Already at preschool age, 16-21% are diagnosed with chronic diseases, which, when entering school, seriously complicate adaptation to the school environment and are the cause of further deterioration in health and academic performance [5,12].

An essential factor determining the state of health of the population is the maintenance of optimal physical activity throughout the life of each citizen [11,12].

Many authors point to a close relationship between indicators of physical development and physical fitness in different population groups [13,14].

The above problems dictate the need to study the degree of physical development and motor quality of children of different ages in the conditions of scientific and technological progress.

Purpose of the study: the purpose of this study is to identify gender, age and constitutional features of physical development and motor qualities in children aged 4-7 years in Bukhara.

Materials and methods

The work was carried out on the basis of 4 preschool institutions Bukhara.

To solve the tasks set, the parameters of physical development and motor qualities of children of the period of the first childhood of 4-7 years were measured, on the basis of 4 preschool institutions in Bukhara. The studies were carried out from September 2020 to May 2022 twice a year, at the beginning (September) and at the end (May) of the school year. Over the past period, 3369 examinations of children aged 4-7 years were carried out, which is 13% of the coverage of all children of this age group in Bukhara, of which 1756 (52.12%) are girls and 1613 (47.88%) are boys.

Anthropometric measurements of the parameters of physical development and functional tests were carried out in the medical office. Motor qualities were tested during physical education classes with the involvement of educators and physical education instructors.

Thus, the following groups of research methods were used: anthropometric, functional. Testing of motor qualities, static.

Results and discussions

The study of the integral indicator of physical development in boys 4-7 years old revealed the largest proportion of them with an average level of development at all ages. At the same time, at the age of 5, there are fewer such children, as in girls, and with levels above the average and high, on the contrary, more (39.8%).

Also, an increase in the proportion of boys with high and above average levels of physical development is noted at 6 and 7 years old (up to 31.1 % and 29.6%, respectively) and slightly - at 4 years old (26.6%).

At the same time, among them, the proportion of boys with low and below average levels decreases to 20.2% at 4 years old, to 19.2% at 6 years old, and to 17.2% at 7 years old, while with an average level it decreases, is about 50%.

Motor qualities

Testing the speed-strength endurance of the trunk muscles - "Lifting the trunk up to 30 seconds" - showed a significant ($P < 0.01$) annual improvement in results from 4 to 7 years. The annual increase has maximum values at 5 and 6 years (up to 2.4% per year), and by the age of 7 it decreases (up to 2.2 times per year).

The dynamic strength of the muscles of the lower extremities - "Bum in length from the spot" annually significantly ($P < 0.01$) changes incrementally from 4 to 7 years. The maximum annual increase in the values of the indicator falls on the age of 5 years -19.4 cm per year, then it gradually decreases to 13.2 (6 years) and 6.8 (7 years) cm per year.

Table 1

Distribution of boys by levels of physical development (in %)

Age/Level	Short	Belowtheaverage	Average	Aboveaverage	High
4 oftheyear	8,6	12,0	54	12,1	16,0
5 years	5,9	13,0	43	26,1	17,2
6 years	7,1	15,0	52	23	9,0
7 years	5,4	12,4	52	18,1	13,1

The values of the indicator of speed - "Running for 30 meters on the move" significantly ($P < 0.01$) also change from 4 to 7 years. The value of growth annually decreases from 5 years (1.8'sec), which indicates a deterioration in this indicator in boys.

In the test for active flexibility of the spine and hip joints - "Seated forward bend", a significant ($P < 0.01$) deterioration in the values of the indicator occurs only from 4 to 5 years. They continue to worsen, but not reliably. The annual increase in test results (Fig. 3) has negative values throughout, increasing from the age of 5, and its maximum values are determined at the age of 6-7 years (0.6 cm per year).

Most motor skills improve at the fastest rate at 5 years, with the exception of active spinal flexibility. The next thing to note is that at 6-7 years old motor (except for the dynamic strength of the muscles of the lower extremities at 7 years old). At the same time, the improvement in speed-strength endurance of the muscles of the body is prolonged for all ages (tab. 2)

Table 2

Annual increase in the indicator of physical development and motor qualities in boys aged 5-7 years in Bukhara

Sign	5 years	6 years	«> 7 years
Bodylength, cm			
Bodyweight, kg			
Chestcircumference, cm			
Shoulder circumference at rest, cm			
Vitallungcapacity, ml			
Martin-Kushelevskyttest, sec			
Dynamometry of the right hand, kg			
Dynamometry of the left hand, kg			
Torso lift 30 sec q.va times			
Standinglongjump, cm			
Run for 30 m immediately, sec			
Forwardbend, sitting, cm			

Highlight periods in white are low values of annual growth, green are high, and red a maximum. When analyzing the integral indicator in boys, it showed that in all ages, most of them have an average level of development of motor qualities (Table 3).

If the proportion of such children is 50% at 5-7 years old, then at 4 years old it increases, mainly due to a decrease with levels below the average (up to 13.5%). At 5 and 7 years old, the proportion of boys with a level of motor qualities above medium and high (up to 32.1% and 34%, respectively), and at 6 years old - slightly (up to 2(pJ%)), while their proportion with a level below the average and low at these ages

Table 2

Distribution of boys by the level of motor qualities (in %)

Age	Short	Belowtheaverage	Average	Aboveaverage	High
4 oftheYear	7,9	6,1	65,0	16,2	6,9
5 years	4,8	14,0	51,1	21,1	12,1
6 years	6,9	13,8	52,2	15,0	12,2
7 years	3,6	14,4	49,1	16,9	16,3

Based on the foregoing, it can be concluded that the period of intensive improvement in motor qualities in most indicators in boys 4-7 years old coincides in time with the maximum increase in body length, vital capacity of the lung and strength of the muscles of the hands at 5 years old (tab. 2). However, at 6 years old their increase in most indicators of physical development and motor qualities is quite high, except for the test of Martine Kushelevsky and Carpal dynamometry. By the age of 7, it again reaches its maximum values at the circumference of the shoulder, however, at the length of the bodies, the vital capacity of the lungs (VC) and the dynamic strength of the lower extremities decreases. At the same time, at the age of 5, the proportion of boys with high and above average levels of both physical development and motor qualities increases among them (tab.2,3).

Conclusions

1. In children of the period of first childhood (4-7 years old), gender differences in physical development are detected from 5-7 years old, and motor qualities - from 4-5 years old. and motor qualities in girls - at 6 years old and in boys - at 5 and 7 years old.
2. The predominance of the average monthly increase in indicators of physical development (by 2,2-35 times) and motor qualities by (4.8-37 times), regardless of gender, is observed in children after summer holidays, while over the rest of the year its intensity decreases ($P < 0.05$).
3. The constitutional features of the physical development of children aged 4-7 years are characterized by significantly ($P < 0.05$) large values of indicators in the macrosomal somatotype, minimal in the microsomal and medium in the mesosomal. The constitutional features of motor qualities are characterized by the predominant development of speed - and active flexibility of the spine and hip joints in children of the macrosomal somatotype, the dynamic strength of the muscles of the lower extremities - mesosomal and speed - strength endurance of the trunk muscles - microsomal.
4. In children of the period of first childhood (4-7 years), the usual variant of the development of independence from sex is found in 64.5%, shortened - in 18.4% and extended - in 17.1%. Age-related features are characterized by an increase in the proportion of boys with a shortened (in 8, 7 times) and extended (2.2 times) developmental variants from 4 to 5 years ($P < 0.05$).
5. In children of the period of first childhood (4-7 years) gender differences in the component composition of the body are characterized by the predominance of relative fat mass in girls (by 5.8%) and muscle mass in boys (by 9.3%). no differences were found in children ($P < 0.05$).

6. Differences in the proportional level of variation in children of the first childhood period (4-7) are characterized by the predominance of the macromembrane type (80.0% in girls and 78.6% in boys), less frequency of occurrence of the mesomembral type (9.8% in girls and 19.1% in boys) and micromembrane (10.22% in girls and 2.3% in boys), and micromembrane types from 4 to 5 years, and in girls by 4.3-18 times from 5 to 6 years ($P < 0.05$).

LIST OF REFERENCES:

1. Sharipova G.I. Paediatric Laser Dentistry // *International Journal of Culture and Modernity* ISSN: 2697-2131., 2022; 12(Jan):33-379.
2. Sharipova G.I. Light and laser radiation in medicine // *European journal of modern medicine and practice* 2022; 2(1):31-36 EJMMP ISSN: 2795-921X.-P.
3. Sharipova Gulniho Ildiyevna. The effectiveness of the use of magnetic-infrared-laser therapy in traumatic injuries of oral tissues in preschool children // *Academic leadership*. 2020; 21(1): ISSN 1533-1812
4. Sharipova Gulniho Ildiyevna. Discussion of results of personal studies in the use of mil therapy in the treatment of trauma to the oral mucosa // *European Journal of Molecular medicine* 2022; 2(2):150-157 March Published by eJournals PVT LTDDOI prefix: 10.52325 Issued Bimonthly Requirements for the authors.
- a. Бауэр В.Г. Социальная значимость физической культуры и спорта в современных условиях развития России // *Теория и практика физической культуры* 2001; 1:50-56.
5. Губа В.П. и др. Оценка физического развития детей по данным морфологических наблюдений / В.П. Губа, Я.С. Татаринцев // *Теория и практика физ. культуры* 2003; 208.
6. Индреев М.Х. Мониторинг физического состояния и актуальность и организации в раннем возрасте / М.Х. Индреев, М.М. Киржинов // *Оздоровление нации и формирование здорового образа жизни насел. Материалы III всероссийской научно-практической Конференция*. - Нальчик: ВНИИФК и С. 2007; 54-55.
7. Кузмин А.Ф. Здоровье и умственная работоспособности первоклассников в период адаптации к школьному обучению / А.Ф. Кузмин, Н.В. Тихомиров // *IV Сибирский физиологический съезд: тез. докл.* - Барнаул, 2008; 137.
8. Косимов Х.О., Манасова И.С., Жумаева А.А. Болалар ва ўсмирлар гигиенаси. / Ўқув қўлланма. Бухара, 2019; 270.
9. Жаворников Б.Д. и др. Физкультурно-оздоровительная работа в субъектах Российской Федерации / Б.Д. Жаворников, М.В. Тихомиров, Е.А. Савенкова // *Инновационные технологии в развитии современного спорта: материалы международной научной конференции "Восток-Россия-Запад"-Иркутск*, 2008; 22-27.
10. Звягина В.В. Использование оздоравливающих факторов природной среды в организации образовательного процесса / В.В. Звягина // *II Рос. науч. практ. конф – Екатеринбург* 2004; 74-79.
11. Яковенко О.Н. Некоторые результаты динамического наблюдения состояния здоровья студентов-медиков по данным анкетирования / О.Н. Яковенко, Т.С. Прокопчук // *Инновационный форум: матер. Кругл ого стола форума - выставки*. - Иркутск, 2008; 97-108.
12. Зайцев А.А. Изменение выраженности жировой массы у спортсменок 9-20 лет различных соматических типов // *Вести ВГУ* 2005; 2:126-130.
13. Шарайкина Е.П. Морфофункциональные и биохимические характеристика молодых людей Красноярска // *Актуальные проблемы морфологический научные труды – Красноярск* 2005; 227-229.

Entered 20.02.2023