



New Day in Medicine
Новый День в Медицине

NDM



TIBBIYOTDA YANGI KUN

Ilmiy referativ, marifiy-ma'naviy jurnal



AVICENNA-MED.UZ



ISSN 2181-712X.
EiSSN 2181-2187

6 (56) 2023

Сопредседатели редакционной коллегии:

**Ш. Ж. ТЕШАЕВ,
А. Ш. РЕВИШВИЛИ**

Ред. коллегия:

М.И. АБДУЛЛАЕВ
А.А. АБДУМАЖИДОВ
А.Ш. АБДУМАЖИДОВ
Р.Б. АБДУЛЛАЕВ
Л.М. АБДУЛЛАЕВА
М.М. АКБАРОВ
Х.А. АКИЛОВ
М.М. АЛИЕВ
С.Ж. АМИНОВ
Ш.Э. АМОНОВ
Ш.М. АХМЕДОВ
Ю.М. АХМЕДОВ
Т.А. АСКАРОВ
М.А. АРТИКОВА
Ж.Б. БЕКНАЗАРОВ (главный редактор)
Е.А. БЕРДИЕВ
Б.Т. БУЗРУКОВ
Р.К. ДАДАБАЕВА
М.Н. ДАМИНОВА
К.А. ДЕХКОНОВ
Э.С. ДЖУМАБАЕВ
Н.Н. ЗОЛОТОВА
А.Ш. ИНОЯТОВ
С. ИНДАМИНОВ
А.И. ИСКАНДАРОВ
С.И. ИСМОИЛОВ
Э.Э. КОБИЛОВ
Д.М. МУСАЕВА
Т.С. МУСАЕВ
Ф.Г. НАЗИРОВ
Н.А. НУРАЛИЕВА
Б.Т. РАХИМОВ
Х.А. РАСУЛОВ
Ш.И. РУЗИЕВ
С.А. РУЗИБОЕВ
С.А. ГАФФОРОВ
С.Т. ШАТМАНОВ (Кыргызстан)
Ж.Б. САТТАРОВ
Б.Б. САФОЕВ (отв. редактор)
И.А. САТИВАЛДИЕВА
Д.И. ТУКСАНОВА
М.М. ТАДЖИЕВ
А.Ж. ХАМРАЕВ
А.М. ШАМСИЕВ
А.К. ШАДМАНОВ
Н.Ж. ЭРМАТОВ
Б.Б. ЕРГАШЕВ
Н.Ш. ЕРГАШЕВ
И.Р. ЮЛДАШЕВ
Д.Х. ЮЛДАШЕВА
А.С. ЮСУПОВ
М.Ш. ХАКИМОВ
Д.О. ИВАНОВ (Россия)
К.А. ЕГЕЗАРЯН (Россия)
DONG JINCHENG (Китай)
КУЗАКОВ В.Е. (Россия)
Я. МЕЙЕРНИК (Словакия)
В.А. МИТИШ (Россия)
В.И. ПРИМАКОВ (Беларусь)
О.В. ПЕШИКОВ (Россия)
А.А. ПОТАПОВ (Россия)
А.А. ТЕПЛОВ (Россия)
Т.Ш. ШАРМАНОВ (Казахстан)
А.А. ЩЕГОЛОВ (Россия)
Prof. Dr. KURBANHAN MUSLUMOV(Azerbaijan)
Prof. Dr. DENIZ UYAK (Germany)

www.bsmi.uz

<https://newdaymedicine.com>

E: ndmuz@mail.ru

Тел: +99890 8061882

**ТИББИЁТДА ЯНГИ КУН
НОВЫЙ ДЕНЬ В МЕДИЦИНЕ
NEW DAY IN MEDICINE**

*Илмий-рефератив, маънавий-маърифий журнал
Научно-реферативный,
духовно-просветительский журнал*

УЧРЕДИТЕЛИ:

**БУХАРСКИЙ ГОСУДАРСТВЕННЫЙ
МЕДИЦИНСКИЙ ИНСТИТУТ
ООО «ТИББИЁТДА ЯНГИ КУН»**

Национальный медицинский
исследовательский центр хирургии имени
А.В. Вишневского является генеральным
научно-практическим
консультантом редакции

Журнал был включен в список журнальных
изданий, рецензируемых Высшей
Аттестационной Комиссией
Республики Узбекистан
(Протокол № 201/03 от 30.12.2013 г.)

РЕДАКЦИОННЫЙ СОВЕТ:

М.М. АБДУРАХМАНОВ (Бухара)
Г.Ж. ЖАРЫЛКАСЫНОВА (Бухара)
А.Ш. ИНОЯТОВ (Ташкент)
Г.А. ИХТИЁРОВА (Бухара)
Ш.И. КАРИМОВ (Ташкент)
У.К. КАЮМОВ (Тошкент)
Ш.И. НАВРУЗОВА (Бухара)
А.А. НОСИРОВ (Ташкент)
А.Р. ОБЛОКУЛОВ (Бухара)
Б.Т. ОДИЛОВА (Ташкент)
Ш.Т. УРАКОВ (Бухара)

6 (56)

2023

июнь

Received: 20.05.2023, Accepted: 30.05.2023, Published: 15.06.2023.

UDC 616.36-002.951.21(611.36):616.89-02-089

TREATMENT OF MULTIPLE LIVER ECHINOCOCCI

Khamdamov B.Z. <https://orcid.org/0000-0003-3569-6688>

Mirkhodjaev I.A. <https://orcid.org/0000-0002-9387-0384>

Akhrorova L.B. <https://orcid.org/0000-0003-3569-6688>

Bukhara State Medical Institute named after Abu Ali ibn Sina Uzbekistan Bukhara, A.Navoi st. 1
Tel: +998(65) 223-00-50 e-mail: info@bsmi.uz

✓ **Resume**

In article results of the operative treatment 67 sick with plural Echinococcus liver. They Are Executed following interference: Ideal Echinocectomy beside 8(17,4%), cistperiectiony beside 6(13%), Echinocectomy beside 24(52,2%)sick. The Remaining cavity liver is processed betadin with exposure 5 ± 0,2; which in 98 % events has brought about ruins scolex parasite. At postoperative period given preparation was entered through drenaj tube in remaining cavity that has allowed to conduct the preventive maintenance of the relapse of the disease. In early and remote postoperative period beside handled sick relapse of the disease did not exist.

Keywords: *echinococcus liver, scolecsizid, betaden, relapse*

ЛЕЧЕНИЕ МНОЖЕСТВЕННЫХ ЭХИНОКОККОВ ПЕЧЕНИ

Хамдамов Б.З. <https://orcid.org/0000-0003-3569-6688>

Мирходжаев И.А. <https://orcid.org/0000-0002-9387-0384>

Ахророва Л.Б. <https://orcid.org/0000-0003-3569-6688>

Бухарский государственный медицинский институт имени Абу Али ибн Сины, Узбекистан,
г. Бухара, ул. А. Навои. 1 Тел: +998 (65) 223-00-50 e-mail: info@bsmi.uz

✓ **Резюме**

В статье проанализированы результаты оперативного лечения 67 больных с множественным и рецидивным эхинококкозом печени. Выполнены следующие вмешательства: идеальная эхинококкэктомия у 8(17,4%), цистперицестэктомия у 6(13%), полуоткрытая эхинококкэктомия у 24(52,2 %) больных. Остаточная полость печени обработана бетадином с экспозицией 5 ± 0,2; который в 98 % случаев привело к гибели сколексов и протесколексов паразита. В послеоперационном периоде данный препарат вводился через дренажную трубку в остаточную полость, что в 21 случаях позволило провести профилактику рецидива заболевания. В раннем и отдаленном послеоперационном периоде у оперированных больных рецидивов заболевания не наблюдалось.

Ключевые слова: *эхинококк, сколексоцид, бетадин, рецидив.*

KO'P SONLI JIGAR EXINOKOKKOZINI DAVOLASH

Khamdamov B.Z. <https://orcid.org/0000-0003-3569-6688>

Mirkhodjaev I.A. <https://orcid.org/0000-0002-9387-0384>

Akhrorova L.B. <https://orcid.org/0000-0003-3569-6688>

Abu Ali ibn Sino nomidagi Buxoro davlat tibbiyot instituti, O'zbekiston, Buxoro, st. A. Navoiy. 1
Tel: +998 (65) 223-00-50 e-mail: info@bsmi.uz

✓ Rezyume

Maqolada ko'p va takroriy jigar exinokokkozi bilan og'rigan 67 nafar bemorni jarrohlid davolash natijalari tahlil qilingan. Quyidagi tadbirlar amalgal oshirildi: 8(17,4%) bemorda ideal echinokokkektomiya, 6(13%) bemorda sistoperitsistekomiya, 24(52,2%) bemorda yarim yopiqexinokokkektomiya. Jigarning qoldiq bo'shilg'i $5 \pm 0,2$ ta'sir qilish bilan betadin bilan davolandı; bu 98% hollarda parazitning skoleks va protekoleks o'limga olib keldi. Operatsiyadan keying davrda ushbu preparat drenaj trubkasi orqali qoldiq bo'shliqqa kiritildi, bu 21 holatda kasallikning qaytalanishini oldini olishga imkon berdi. Operatsiya qilingan bemorlarda operatsiyadan keying erta va kechki davrda kasallikning qaytalanishi kuzatilmadi.

Kalit so'zlar: exinokokk, skoleksotsid, betadin, relaps.

Relevance

The most radical and modern method of treating patients with liver echinococcosis (EP) is surgical [4]. The traditionally performed method of open echinococcectomy does not fully satisfy surgeons, because the frequency of developing postoperative complications remains high, reaching from 35% to 40% - [2]. Multiple lesions of liver echinococcal cysts (IEL) occur in 6.4-28% of cases [3], more often recurrence occurs precisely in the share of multiple forms of EP.

Purpose of the study. The purpose of this study is to study the results of surgical treatment of multiple and recurrent forms of EP, to develop ways to improve the results.

Material and methods

The results of surgical treatment of 67 patients with multiple EP were analyzed. In 46 patients with primary multiple EP, localization of cysts in the right lobe of the liver was observed in 20 (43%) patients, in the left lobe of the liver in 8 (17%), in both lobes of the liver in 21 patients. Uncomplicated forms of MEP occurred in 28 patients, complicated forms in 18 patients. In the preoperative period, the following complications were observed: cyst rupture into the biliary tract in 4 (22%), rupture into the abdominal cavity in 1 (5.6%), rupture into the pleural cavity in 2 (11%), compression of the biliary tract with the development of a mechanical jaundice in 5 (28%), cyst calcification in 2 (11%), cyst suppuration in 4 (22%) patients. With multiple liver echinococcosis, the following surgical interventions were performed: ideal echinococcectomy in 8 (17.4%), cystpericistectomy in 8 (17.4%), echinococcectomy + cystpericistectomy in 6 (13%), semi-closed echinococcectomy in 24 (52.2%) sick. The largest percentage of operations were 23 patients with external drainage of the residual cavity, and in 17 cases, options for reducing residual cavities were used (partial pericystectomy, invagination, capitonnage). The residual cavity of the liver was treated with betadine, which has a detrimental effect on the germinal elements of the parasite. In 100 mg of betadine, consisting of active iodine in an amount of 10 mg, when treating the residual echinococcal cavity with an exposure of 5 ± 0.2 , in 98% of cases it provides a reliable disinfecting effect on all elements of the echinococcal cyst. Unlike many other antiscolexicidal drugs used in the clinical practice of liver echinococcosis surgery, which are hepatotoxic, betadine is free from such disadvantages. Using in the postoperative period the introduction of betadine through a drainage tube into the residual cavity of the liver allowed to minimize the cases of suppuration of the residual cavity, and served as a reliable germicide in the prevention of recurrence of the disease. In 4 cases, with complicated EP, a breakthrough of the contents of the cyst into the biliary tract occurred. These patients underwent endoscopic papillosphincterotomy (EPST), sanitation of the extrahepatic biliary tract with decasan + open liver echinococcectomy with drainage of the residual cavity. In the postoperative period held antiparasitic and hepatotropic therapy. After the operation of traditional liver echinococcectomy in the early postoperative period, the following complications developed: exudative pleurisy in 3 (6.5%) cases, an allergic reaction in 2 (4.3%), and the formation of a biliary fistula in 6 (13%) patients. In the diagnosis of exudative pleurisy, along with polypositional X-ray examination, ultrasound of the pleural cavity is of particular importance. The advantage of the ultrasound method is that in the pleural cavity, the accumulation of fluid with a volume of about 10 ml is clearly visualized by this method, while with fluoroscopy, the fluid in the pleural cavity is detected only at a volume of more than 100 ml. Under the control of ultrasound, it is possible to determine the depth, localization and the possibility of repeated punctures without damaging the internal organs.



All three patients with reactive pleurisy underwent daily puncture of the pleural cavity, followed by the use of a broad-spectrum antibiotic for 5-7 days. The criterion for the effectiveness of puncture sanitation of the cavity was the elimination of manifestations of intoxication and the absence of accumulation of exudate in the pleural cavity. Bile leakage during external drainage of residual liver cavities was observed in 6 patients, which stopped by 15-16 days after conservative treatment. In the postoperative period, software dynamic ultrasound was performed to control the regression of the residual liver cavities. Complete reduction of residual cavities in operated patients occurred by 26 ± 5.6 days. In 6 cases, after surgery for MEP, a relapse of the disease was observed, which was successfully cured by puncture-aspiration echinococcectomy under ultrasonography control.

After traditional and minimally invasive methods of MEP treatment, 3 courses of antiparasitic therapy with albendazole (10-12 mg/kg of the patient's weight) were carried out for 28 days with a break interval of 1 month. Of the 21 recurrent forms of MEP, cysts were localized mainly in segments VII, VIII of the liver. The diagnosis was established on the basis of ultrasound, CT, MRI and high titers antibodies - 400 or more, enzyme immunoassay (ELISA). All patients underwent percutaneous minimally invasive surgical interventions – minilaparotomy. All patients underwent percutaneous minimally invasive surgical interventions – minilaparotomy. Indications for such interventions were sonographic pictures of echinococcal cysts II, III types according to H.G. Harbi, suppuration of the parasitic cavities of the liver, 8-10 cm in size, the severity of the general condition of patients. The technique for performing percutaneous interventions from a mini-access under ultrasound control was as follows: all manipulations were performed under intravenous ketanalev anesthesia. Before the intervention, an ultrasound scan was performed, the place for minilaparotomy was determined, observing the principle of antiparasiticity. The cyst or the residual cavity was punctured, the fluid was evacuated, the wound was expanded along the needle, the residual cavity was sanitized and drained with PVC drains. Antiparasitic treatment of the residual cavity was carried out with a solution of betadine with an exposure of 15 minutes. The terms of drainage of the cavities averaged 16.8 ± 4.2 days.

Results and discussions

After minimally invasive interventions under ultrasound control, puncture methods for extracting fluid from the pleural cavity were performed in 2 patients with reactive pleurisy. No other complications were observed in the early and late postoperative period. Patients with recurrent EP in the postoperative period underwent chemotherapy with albendazole at a dose of 10-12 mg/kg per day, recommended by WHO (1996). According to it, patients with a live parasite and a thin-walled cyst (type I and II cyst according to Garbi) started treatment 4 days before surgery and continued for 2 weeks after surgery at a daily dose of up to 200 mg. The drug albendazole with severe fibrosis and calcification of the cyst (III and IV type of cyst according to Garbi) tried to start treatment earlier and increased the preoperative daily dose of albendazole to 300 mg, and in the postoperative period, the intake of this drug remained unchanged.

In case of intraoperative difficulties and doubts about the observance of the principle of aparasiticity of the stages of the operation, the chemotherapeutic postoperative daily dose of albendazole was increased to 300 mg and the course of treatment was extended to 1 month. No complications of chemotherapy were observed. The analysis of the results of surgical treatment of liver echinococcosis in patients showed that intraoperative complications are not so significant (4.2%). Clear advantages of external drainage of an unreduced fibrous cavity over the option of drainage with a reduced residual cavity were noted - omentoplasty, capitonnage and invagination methods.

Conclusion

Thus, the most effective methods of surgical treatment of liver echinococcosis should be considered cystpericistectomy, ideal echinococcectomy, in which the risk of recurrence of the disease is reduced to zero, and from open methods, echinococcectomy is external drainage of the residual cavity with repeated treatment of the residual cavity with betadine. Echinococcectomy of the liver from a mini-access in case of relapse of the disease creates favorable conditions for compliance with the principles of aparasitism and antiparasitism, being a low-traumatic operation, strictly observing the principles of aparasitism and antiparasitism, while the time spent by the patient in the hospital is rarely reduced, the development of postoperative complications was not observed.

LIST OF REFERENCES:

1. Мирходжаев И.А, Ахророва Л.Б. Совершенствование методов хирургического лечения множественных и рецидивных форм эхинококкоза печени Новый День Медицины (2023) 2(52):270-273 <https://newdaymedicine.com/index.php/2023/02/26/>
2. Ахророва Л.Б. Результаты хирургического лечения больных острым гангренознекротическим парапроктитом Вопросы науки и образования (2020) 29(154):85-90.
3. Ахророва Л.Б. Выбор хирургической тактики лечения больных острым парапроктитом Вопросы науки и образования, (2021) 27(152):51-57.
4. Газиев К.У. и др. Ампутации на уровне голени при критической ишемии у больных сахарным диабетом Биология и интегративная медицина, (2021) 1(48).
5. Хамдамов Б.З. Метод лазерной фотодинамической терапии в лечении раневой инфекции при синдроме диабетической стопы Проблемы биологии и медицины, (2020) 1:142-148.
6. Davlatov S.S. et al. Plasmapheresis in the treatment of cholemic endotoxicosis Academic Journal of Western Siberia, (2013) 9(1):30-31.
7. Hamdamov B.Z. Optimization of methods of local treatment of purulent-necrotic lesions of the foot in diabetes mellitus New day in medicine, (2018) 4:24.
8. Khamdamov B.Z. et al. Method of prevention of postoperative complications of surgical treatment of diabetic foot syndrome European science review (2018) 9-10(2):194-196.
9. Khamdamov B.Z. et al. The role and place laser photodynamic therapy in prevention postoperative complication at treatment of diabetic foot syndrome Applied Sciences: challenges and solutions (2015):27-31.
10. Rustamov M.I., Davlatov S.S., Saydullaev Z.Y., Rustamov I.M. Choice of surgical tactics of treatment of patients with acute paraproctitis. Journal of hepatogastroenterology research, (2020) 2(1):26-29.
11. Rustamov M.I., Davlatov S.S., Saydullaev Z.Y., Rustamov I.M. Results of surgical treatment of patients with acute gangrenous - necrotic paraproctitis. Journal of hepatogastroenterology research, 2020.
12. Ахметов Д.Г. Радикальные операции при гидатидозном эхинококкозе печени - Анналы хирургической гепатологии (2007) 12(3):41.
13. Гончаров А.Б, Коваленко Ю.А, Икромов Р.З. “Сложний” эхинококкоз печени Ж.Анналы хирургический гепатологии” (2021) 26(4):32-40.
14. Ахметов Д.Г. Совершенствование оперативных доступов при гидатидозном эхинококке печени - Вестник Киргизского университета (2007) 7(7):150-153.
15. Шангареева Р.Х. Хирургическое лечение множественного эхинококкоза печени у детей 2005.
16. Киртанасов Я.П. Чрезкожные вмешательства в лечении больных многокамерных гидатидным эхинококкозом печени Ж. Вестник новых медицинских технологий (2019) 2:23-32.
17. Панфилов К.А., Иванов С.А. Выбор способа и анализ результатов применения малоинвазивных методов лечения гидатозного эхинококкоза печени Ж. Вестник экспериментальной и клинической хирургии (2019) 11(4):222-229.
18. Мирходжаев И.А., Комилов С.О. Прогностическое значение цитокинового профиля при множественном эхинококкозе живых организмов Новый день в медицине 2020 4(32):730-732 <https://newdaymedicine.com/index.php/2021/01/18/206-4-32-4-2020>

Entered 20.05.2023

