



New Day in Medicine
Новый День в Медицине

NDM



TIBBIOVIYOTDA YANGI KUN

Ilmiy referativ, marifiy-ma'naviy jurnal



AVICENNA-MED.UZ



ISSN 2181-712X.
EiSSN 2181-2187

9 (59) 2023

Сопредседатели редакционной коллегии:

**Ш. Ж. ТЕШАЕВ,
А. Ш. РЕВИШВИЛИ**

Ред. коллегия:

М.И. АБДУЛЛАЕВ
А.А. АБДУМАЖИДОВ
А.Ш. АБДУМАЖИДОВ
Р.Б. АБДУЛЛАЕВ
Л.М. АБДУЛЛАЕВА
М.М. АКБАРОВ
Х.А. АКИЛОВ
М.М. АЛИЕВ
С.Ж. АМИНОВ
Ш.Э. АМОНОВ
Ш.М. АХМЕДОВ
Ю.М. АХМЕДОВ
Т.А. АСКАРОВ
М.А. АРТИКОВА
Ж.Б. БЕКНАЗАРОВ (главный редактор)
Е.А. БЕРДИЕВ
Б.Т. БУЗРУКОВ
Р.К. ДАДАБАЕВА
М.Н. ДАМИНОВА
К.А. ДЕХКОНОВ
Э.С. ДЖУМАБАЕВ
Н.Н. ЗОЛотова
А.Ш. ИНОЯТОВ
С. ИНДАМИНОВ
А.И. ИСКАНДАРОВ
Э.Э. КОБИЛОВ
Д.М. МУСАЕВА
Т.С. МУСАЕВ
Ф.Г. НАЗИРОВ
Н.А. НУРАЛИЕВА
Б.Т. РАХИМОВ
Х.А. РАСУЛОВ
Ш.И. РУЗИЕВ
С.А. РУЗИБОВЕВ
С.А. ГАФФОРОВ
С.Т. ШАТМАНОВ (Кыргызстан)
Ж.Б. САТТАРОВ
Б.Б. САФОВЕВ (отв. редактор)
И.А. САТИВАЛДИЕВА
Д.И. ТУКСАНОВА
М.М. ТАДЖИЕВ
А.Ж. ХАМРАЕВ
А.М. ШАМСИЕВ
А.К. ШАДМАНОВ
Н.Ж. ЭРМАТОВ
Б.Б. ЕРГАШЕВ
Н.Ш. ЕРГАШЕВ
И.Р. ЮЛДАШЕВ
Д.Х. ЮЛДАШЕВА
А.С. ЮСУПОВ
М.Ш. ХАКИМОВ
Д.О. ИВАНОВ (Россия)
К.А. ЕГЕЗАРЯН (Россия)
DONG JINCHENG (Китай)
КУЗАКОВ В.Е. (Россия)
Я. МЕЙЕРНИК (Словакия)
В.А. МИТИШ (Россия)
В.И. ПРИМАКОВ (Беларусь)
О.В. ПЕШИКОВ (Россия)
А.А. ПОТАПОВ (Россия)
А.А. ТЕПЛОВ (Россия)
Т.Ш. ШАРМАНОВ (Казахстан)
А.А. ЩЕГОЛОВ (Россия)
Prof. Dr. KURBANHAN MUSLUMOV (Azerbaijan)
Prof. Dr. DENIZ UYAK (Germany)

www.bsmi.uz

<https://newdaymedicine.com>

E: ndmuz@mail.ru

Тел: +99890 8061882

**ТИББИЁТДА ЯНГИ КУН
НОВЫЙ ДЕНЬ В МЕДИЦИНЕ
NEW DAY IN MEDICINE**

Илмий-рефератив, маънавий-маърифий журнал

Научно-реферативный,

духовно-просветительский журнал

УЧРЕДИТЕЛИ:

**БУХАРСКИЙ ГОСУДАРСТВЕННЫЙ
МЕДИЦИНСКИЙ ИНСТИТУТ
ООО «ТИББИЁТДА ЯНГИ КУН»**

Национальный медицинский
исследовательский центр хирургии имени
А.В. Вишневского является генеральным
научно-практическим
консультантом редакции

Журнал был включен в список журнальных
изданий, рецензируемых Высшей
Аттестационной Комиссией
Республики Узбекистан
(Протокол № 201/03 от 30.12.2013 г.)

РЕДАКЦИОННЫЙ СОВЕТ:

М.М. АБДУРАХМАНОВ (Бухара)
Г.Ж. ЖАРЫЛКАСЫНОВА (Бухара)
А.Ш. ИНОЯТОВ (Ташкент)
Г.А. ИХТИЁРОВА (Бухара)
Ш.И. КАРИМОВ (Ташкент)
У.К. КАЮМОВ (Тошкент)
Ш.И. НАВРУЗОВА (Бухара)
А.А. НОСИРОВ (Ташкент)
А.Р. ОБЛОКУЛОВ (Бухара)
Б.Т. ОДИЛОВА (Ташкент)
Ш.Т. УРАКОВ (Бухара)

9 (59)

2023

сентябрь

UDK 616(072)+616-091(072)-053.3

FREQUENCY OF NEONATAL ATELECTASIS OF THE LUNG AND MORPHOLOGICAL CHANGES

Ruzieva Z.I. Email: G.jurayeva69@gmail.com

Tashkent Medical Academy 100109, Tashkent, Uzbekistan Farabi Street 2. Tel: +99878 1507825;
E-mail: info@tma.uz

✓ Resume

The aim of this study was to address the problems of lung tissue changes in atelectasis. Primary atelectasis of the lungs "enters the respiratory distress syndrome and appears as a separate nosological unit. In this article, specific pathomorphological signs of primary lung atelectasis are studied. As a material, the lungs of babies who died of atelectasis in the neonatal period were studied microscopically. In the results of the microscopic examination, it was found that the lung tissue has an underdeveloped appearance at first glance.

It is observed that the tissue between the alveoli consists of dense tissue and cellular tufts, the blood vessels are wide and full, and it has a structure around which blood clots have appeared. Macrophages, neutrophils, and migrated alveolocytes are detected in the alveolar space. After 7-10 days, it is determined that alterative-proliferative processes escalated and turned into atelectatic pneumonia. As a result, pneumosclerosis, bronchiectasis, and the transformation of bronchi into retention cysts are observed. Often, at the site of atelectasis, connective tissue grows and sclerosing develops.

Keywords: infant, lung, distress syndrome, primary atelectasis newborn infants, atelectasis, respiratory failure syndrome.

ЧАСТОТА ВСТРЕЧАЕМОСТИ АТЕЛЕКТАЗА ЛЕГКОГО У НОВОРОЖДЕННЫХ И МОРФОЛОГИЧЕСКИЕ ИЗМЕНЕНИЯ

З.И. Рузиева Email: G.jurayeva69@gmail.com

Ташкентская Медицинская Академия (ТМА) Узбекистан, 100109, Ташкент, Алмазарский район, ул. Фароби 2, тел: +99878 1507825, E-mail: info@tma.uz

✓ Резюме

Целью данного исследования была, решение проблем изменения ткани легкого при ателектазе. Первичный ателектаз легких «входит в респираторный дистресс-синдром и выступает отдельной нозологической единицей. В статье изучены специфические патоморфологические признаки первичного ателектаза легкого. В качестве материала микроскопически исследовали легкие детей, погибших от ателектаза в период новорожденности. В результате микроскопического исследования установлено, что легочная ткань на первый взгляд имеет недоразвитый вид.

Установлено, что ткань между альвеолами состоит из плотной ткани и клеточных пучков, кровеносные сосуды широкие и полноценные, имеют структуру, вокруг которой появились тромбы. В альвеолярном пространстве обнаружены макрофаги, нейтрофилы и мигрировавшие альвеолоциты. Через 7-10 дней было выявлено, что альтеративно-пролиферативные процессы обострились и переросли в ателектатическую пневмонию. В результате наблюдалось пневмосклероз, бронхоэктатическая болезнь и трансформация бронхов в ретенционные кисты. Нередко на месте ателектаза разрасталось соединительная ткань и развилось склерозирование.

Ключевые слова: новорожденные дети, ателектаз, синдром дыхательной недостаточности.

ЧАҚАЛОҚЛАРДА ЎПКА АТЕЛЕКТАЗИНИНГ УЧРАШ ДАРАЖАСИ ВА МОРФОЛОГИК ЎЗГАРИШЛАРИ

З.И. Рўзиева Email: G.jurayeva69@gmail.com

Тошкент тиббиёт академияси. Ўзбекистон, 100109, Тошкент, Олмазор тумани, Фароби кўчаси 2, тел: +99878 1507825, E-mail: info@tma.uz

✓ Резюме

Ушбу тадқиқотнинг мақсади ателектазда ўпка тўқималарининг ўзгариши муаммосини ҳал қилиш эди.

Ўпканинг бирламчи ателектази “респиратор дистресс синдромга кириб, алоҳида нозологик бирлик кўринишида учрайди. Ушбу мақолада ўпка бирламчи ателектазининг ўзига хос патоморфологик белгилари ўрганилган. Материал сифатида неонатал даврда ателектаздан ўлган чақалоқлар ўпкаси микроскопик жиҳатдан ўрганилди. Микроскопик текширув натижаларида ўпка тўқимаси бир қарашда чала ривожланган кўринишга эгалиги аниқланди.

Альвеолалар оралиги тўқимаси бир-бири билан қўшилиб кетган зич ҳолдаги тўқима ва ҳужайрали тутамлардан иборатлиги, қон томирлари кенг ва тўлақонли, атрофига қон қуйилишлар пайдо қилган тузилишга эгалиги кузатилади. Бирламчи ателектаз чақалоқлар ўлиmidан 2-3 кун олдин ривожланган бўлса, ўпка тўқимасида яллигланиш ривожланганлиги кузатилади, яъни альвеолалар бўйлигида макрофаглар, нейтрофиллар, кўчиб тушган альвеолоцитлар аниқланади. 7-10 кун ўтгандан кейин альтератив-пролифератив жараёнлар авж олиб ателектатик пневмонияга айланганлиги аниқланади. Унинг оқибатида пневмосклероз, бронхоэктазлар ва бронхларнинг ретенцион кистага айланиши кузатилади. Кўпинча ателектаз ўрнида бириктирувчи тўқима ўсиб, склерозланиш ривожланади.

Калит сўзлар: янги туғилган чақалоқлар, ателектаз, нафас олиш етишмовчилиги синдроми.

Relevance

In the early neonatal period, pulmonary atelectasis is a manifestation of respiratory failure syndrome (RFS), which often leads to acute respiratory failure. Lung atelectasis in infants is a primary failure to open lung alveoli or alveolar collapse in the first two days of life due to the specific structure of the lungs and insufficient development of the central system of respiratory regulation [1, 4, 7, 9, 10, 11, 17]. Infantile atelectasis is a part of the "Respiratory Failure Syndrome". The relevance of this problem for paediatrics is due to the large number of causes of lung tissue deterioration in children of one month old. According to the World Health Organization (WHO), its incidence averages 1% of all newborns, and pulmonary atelectasis occurs in up to 14% of cases in premature infants with a body weight of less than 2500 g. Atelectasis occurs in 65% of babies born before 30 weeks gestation who did not receive prenatal prophylaxis with steroid hormones, 35% of those who received prenatal prophylaxis, 25% and 10% of those born at 30-34 weeks, respectively [3,5,6,12,16]. RFS, including surfactant deficiency as a risk factor for the development of pulmonary atelectasis and all conditions leading to premature lung development, including: neonatal and fetal asphyxia, morphofunctional failure, impaired cardiopulmonary adaptation, pulmonary hypertension, metabolic disorders, i.e., acidosis, hypertension, metabolic disorders, etc. [5,5,6,12,16] i.e. acidosis, hypoproteinaemia, hypofermentosis, electrolyte metabolism disorders; diabetes mellitus, haemorrhage in pregnant women, caesarean section, second births of twins.

The aim of the research

Analysis of risk factors for pulmonary atelectasis in infants, determination of pathomorphological changes developing in the focus of atelectasis depending on pregnancy age, life after birth, and evaluation of diagnostic value in "respiratory failure syndrome".

Materials and methods

Autopsy data of 52 preterm and 34 preterm infants who died of pulmonary insufficiency during 2020-2022 were obtained as material. First, the medical history and autopsy report of the infants were analysed. The preterm infants were divided into the following groups according to weeks of gestation: Group 1, 12 (23.1%) 22-27 weeks; Group 2, 18 (34.6%) 28-32 weeks; Group 3, 22 (42.3%) babies born at 33-37 weeks who died of pulmonary insufficiency (Table 1). Of these, group 1, i.e., 22-27-week-old, profoundly premature infants with very low birth weight who died within 1 hour of birth, was taken as a control group. The aim was to identify the first morphological changes in the lungs of this group of children that led to the formation of atelectasis. The main cause of death in this group was postnatal asphyxia. Prenatal and intrapartum risk factors from the mother's side, pathologies from the placenta and umbilical cord were identified in all cases. Anthropometric parameters of the examined children are presented in Table 1.

Table 1

Anthropometric indicators of premature infants, $M \pm m$

№	Group of premature infants	N	Time of pregnancy	Body weight, g	Height, cm
1	22-27 weeks	12	25,2 $\pm$ 0,4	654 $\pm$ 24,3	29,5 $\pm$ 1,6
2	28-32-weeks	18	29,8 $\pm$ 0,6*	1067 $\pm$ 84,7*	38,4 $\pm$ 4,6*
3	33-37-weeks	22	35,3 $\pm$ 0,7**	1986 $\pm$ 124,6**	43,2 $\pm$ 8,5**

* - $P \leq 0.05$ - reliability of the difference in pregnancy term, body weight and height in group 2 compared to group 1.

** - $P \leq 0.05$ - reliability of the difference in pregnancy term, body weight and height in group 3 compared to group 1.

According to the results of clinical and anamnestic analysis, the shortest-lived children and increasing cardiorespiratory failure as a cause of death were observed in the first group of children, and those who lived relatively long - in the 3rd group.

Pieces from different parts of both lungs were taken for histological examination at pathological autopsy. Macroscopic appearance of the lungs was assessed by examining their appearance at autopsy. Lung sections were fixed in 10% neutral formalin, denatured and embedded in paraffin. Histological sections were stained in hematoxylin-eosin, Periodic Acid - Schiff (PAS) reaction and by Van Gieson. The following morphometric calculations were performed.

Results and discussion

Externally, areas of lung atelectasis look like flesh, lung lobes are significantly smaller than their normal size and differ in greyness compared to the surrounding healthy tissue. Foci of atelectasis have been found to have a small appearance, dark red in colour, resembling foci of infarction.

If primary atelectasis develops 2-3 days before the death of babies, the development of inflammation in the lung tissue is observed. As a result, macrophages, neutrophils and migrated alveolocytes are found in the alveolar space. After 7-14 days it is determined that the alterative-proliferative processes have aggravated and passed into atelectatic pneumonia. As a result, pneumosclerosis, bronchiectasis, transformation of bronchi into retention cysts are observed (Fig. 1). It is often determined that the pulmonary vascular system is restored due to connective tissue overgrowth and sclerosing at the site of atelectasis (Fig. 2). If atelectasis foci develop in the subpleural region of the lung, they develop in the posterior basal segments of the lung, and morphologically, partial recession of alveoli and surrounding pathomorphological changes develop to varying degrees.




Fig. 1. Irregular formation of lung tissue, appearance of various cavities and bronchiectasis. Stain: van Gieson method. Size: 10x40.




Fig. 2. Connective tissue overgrowth in the focus of atelectasis. Stain: van Gieson method. Size: 10x40.

The results of microscopic examination showed that the lung tissue appeared underdeveloped at first glance when examined through a small microscope lens. In the lung tissue only bronchi and bronchioles have tubular structures of various sizes and irregular shape. The reticular appearance of the alveolar tissue is not detectable. Respiratory alveoli and communicating spaces are not identified. Alveolar interstitial tissue consists of densely packed tissue and cell bundles (Fig. 3). In such dense lung tissue, wide and full blood vessels are observed, and it has a structure around which thrombi have formed.

When examined under a large microscope objective, it is determined that the bronchial wall consists of only poorly developed tissue structures. It is determined that only one and a few layers of disorderly epithelial cells make up the bronchiolar wall. It is determined that there is a thin tissue surrounding it consisting of dense young histiocytic and lymphoid cells. However, it is not determined whether an alveolar space has appeared in this delicate and young fibre. Therefore, because the lung tissue is young

and immature, it is determined that the alveoli are undifferentiated and consist of dense tissue in the form of primary atelectasis and without alveolar space (Figure 4). In most cases, it is determined that the lung tissue undergoing primary atelectasis is full blooded and there are massive pulmonary haemorrhages (Fig. 5). It is noted that these haemorrhage foci are dense and irregularly shaped haemorrhages appear in tissue without alveolar space. As a result of blood transfusion, the primary pulmonary atelectasis is further thickened and deformed, resembling the red pulp tissue of the spleen. On microscopic examination, primary atelectasis of alveolar tissue is represented by a dense mass of unformed histiocytic and lymphoid cells. In the tissue with this structure, alveoli are not formed, and in the interalveolar tissue alveolocytes, blood vessels and interstitial tissue also have undifferentiated structure. Blood vessels in such dense and unformed tissue are dilated, full and massively filled with blood in most areas (Fig. 6).




Fig.3. Alveolar interstitial tissue consists of densely packed tissue and cell bundles. Stain: van Gieson method. Size: 10x40.




Fig. 4. Foci of primary atelectasis are defined as dense tissue without alveolar space. Stain: van Gieson method. Size: 10x40.




Fig.5. Lung tissue with primary atelectasis has vascular hyperaemia and massive haemorrhages. Stain: van Gieson method. Size: 10x40.




Fig. 6. In the unformed lung tissue, the vessels are dilated and most areas are massively haemorrhaged. Stain: van Gieson method. Size: 10x40.

In infants who developed primary atelectasis in the lung tissue and died after 2-4 days, microscopic examination of the lung tissue revealed that the respiratory part of the lung and the alveolar space appeared in the form of small fissures. In the structure of the tissue between these spaces, it is determined

that histiocytic cells characteristic of proliferative inflammation are activated and hypertrophied, and a dense proliferative inflammatory infiltrate is formed (Fig.7). Macrophages, giant cells and lymphoid cells appear in this inflammatory infiltrate. In the respiratory bronchioles and alveolar part of the lung, the overgrowth of epithelium, including alveolocytes, and the formation of a number of structures are determined.

After the development of primary atelectasis in the lung tissue of young children who died in 7-10 days, microscopic examination of lung tissue revealed that the alveolar tissue thickens and resembles the spleen tissue without voids. Only in places is determined the appearance of spaces of indeterminate shape, in the space of which migrated epithelial and inflammatory cells. In fact, it is determined that the tissue giving rise to the lung tissue, i.e. the tissue of the alveolar space, is sharply thickened and consists of unformed connective tissue (Fig. 8). The outgrowth of such connective tissue, that is, the appearance of tufts of sclerosis, the outgrowth of histiocytic cells and the appearance of fibrous structures between them. When this condition is stained by the histochemical method, i.e. van Gieson's method, it is observed that the fibrous structures of connective tissue developed in the lung tissue are stained red with picrofuchsin (Fig. 9). Blood vessels in it also proliferated, which was manifested by proliferation and increase in endothelial and pericytic cells forming the wall.




Fig. 7. Hypertrophy of proliferative-inflammatory histiocytes, formation of dense proliferative-inflammatory infiltrate between alveoli. Stain: van Gieson method. Size: 10x40.




Fig. 8. Connective tissue overgrowth in the alveolar tissue, i.e. the appearance of sclerotic strands. Stain: van Gieson method. Size: 10x40.




Fig. 9. Increase in fibrous structures of connective tissue in alveolar tissue. Stain: van Gieson method. Size: 10x40.

Conclusion

1. neonatal pulmonary atelectasis is part of the "respiratory distress syndrome (RDS)" and has an overall incidence of 1% of all newborns and 14% in preterm infants. The relevance of atelectasis to paediatrics is that there are many reasons why alveolar lung tissue deteriorates at one month of age.
2. On microscopic examination, the lung tissue at first glance has an underdeveloped appearance. Only bronchioles and bronchioles of tubular structure of various sizes appear in the lung tissue. The reticular appearance of the alveolar tissue is not detected. Respiratory alveoli and communicating spaces are not identified.
3. It is observed that the tissue between alveoli consists of dense tissue and cell bundles, blood vessels are wide and full blooded, has a structure with blood clots around.
4. If primary atelectasis develops 2-3 days before neonatal death, there is inflammation in the lung tissue, i.e. macrophages, neutrophils, migrated alveolocytes are found in the alveolar cavity.
5. After 7-10 days, it is determined that the alternative-proliferative processes have developed into atelectatic pneumonia. As a result, pneumosclerosis, bronchiectasis, transformation of bronchi into retention cysts are observed. Often at the site of atelectasis connective tissue overgrows and sclerosis develops.

LIST OF REFERENCES:

1. Birkun A.A., Zagogulko A.K., Phuc Phat et al. The state of lung surfactant in their immaturity. Archives of Pathology. 1990;2:10-14.
2. Bogdan I.Y. Respiratory therapy of severe forms of RDS in deep premature babies in the early neonatal period. Dissertation, Candidate of Medical Sciences, M., 2000.
3. Bubnova N.I. Artificial ventilation and bronchopulmonary dysplasia in children with low body weight. - Intensive therapy in obstetrics and neonatology: Theses of reports of the I Congress of obstetricians and gynaecologists of the Turkmen SSR, Ashgabat, 1988;41-43.

4. Vanovskaya I.V. Complications of artificial lung ventilation in children. (Literature review). *Voprosy zashchita maternosti i detstva*. 1996;21(1):75-77.
5. Volodin H.H., Degtyarev D.N. Principles of nursing children with extremely low body weight. // *Issues of obstetrics and gynaecology* 2003;2:64-70
6. Glukhovets B.I. Gaivoronsky I.V. et al. Pathogenetic features of respiratory distress syndrome in newborns with extremely low body weight // *Archives of Pathology*. 2005;67(1.):3-5.
7. Grebennikov V.A. Evaluation of the haemodynamic effect of dopamine in newborns with SDR. *Bulletin of the USSR Academy of Medical Sciences*, 1984;9:77-80.
8. Dementieva G.M., Kuzmina T.B., Baleva L.S. et al. Repeated and chronic bronchopulmonary diseases at an early age in children on artificial ventilation in the neonatal period. *Ros. Bulletin of gynecology and pediatrics*. 1997;1:21-24.
9. Evseev S.F., Kutuyavin A.A., Stolovich M.N. Risk factors for the development of bronchopulmonary dysplasia during artificial lung ventilation in newborns // *Emergency conditions in children: Mat. of VI congress. VI congress. Paediatricians of Russia*. M, 2021;107.
10. Evtyukov G.M. Syndrome of respiratory disorders in newborns // *Problems of human health formation in the perinatal period and in childhood: Collection of scientific works*, ed. by N.P. Shabalov. – St.P: Public. "Olga", 2020;172-178.
11. Kuznetsova N.F., Tuzakina U.B., Charushnikova G.A. Features of phospholipid metabolism in women who gave birth to children with respiratory distress syndrome. - *Voprosy zashchita maternosti i detstva*. 2011;2:21-54.
12. Liubsys A. Immaturity of the lungs: neonatal perspective -development of the lung. *Journal of obstetrics and female diseases*. St.Peterburg. 1999;99.
13. American Academy of Pediatrics.Committee on Fetus and Newborn. Surfactant replacement therapy for respiratory distress syndrome.-*Pediatrics*, 1991;87(6):946-947
14. Dorsi M. Pediatric constitutional therapy // *Kinderarztl. Prex*, 1993;61(1):24-27.
15. Forster R. E. Respiratory physiology and lunglipid metabolism. *Ann. Rev. Physiol.*, 2005;47:751.
16. Gibson R.L., Jackson J.C., Twiggs G.A. Bronchopulmonary dysplasia: survival after prolonged mechanical ventilation. // *Am. J.Dis.Child.*, 1988;42:721-732.

Entered 20.08.2023