

HOMILADORLIK VA TUG'RUQDA QUYYI GENITAL TRAKTI INFEKSIYANING AHAMIYATI

Tuxanova D.I.,

Abu Ali ibn Sino nomidagi Buxoro tibbiyot instituti, Buxoro, O'zbekiston.

✓ *Rezyume*

Bexatar onalik yo'nalishi bo'yicha eng muhim vazifa bu perinatal yo'qotishlarning oldini olishdir. Buning intihosi sifatida homila ko'tara olmaslikning sababi sifatida ijtimoiy, tibbiy va biologik omillarning ona va homila vujudiga ta'siri sifatida ko'rib chiqiladi.

Kalit so'zlar: infektsiya, genital tizim, homiladorlik, tug'ruq.

РОЛЬ ИНФЕКЦИИ НИЖНИХ ПОЛОВЫХ ПУТЕЙ В ИСХОДАХ БЕРЕМЕННОСТИ И РОДОВ

Туксанова Д.И.,

Бухарский медицинский институт имени Абу Али ибн Сино, Бухара, Узбекистан.

✓ *Резюме*

Одним из главных направлений в решении проблемы безопасного материнства связано со снижением репродуктивных потерь, которые рассматриваются как конечный результат влияния социальных, медицинских и биологических факторов на здоровье беременной женщины, состояние плода и новорожденного.

Ключевые слова: инфекция, половая система, беременность, осложнения.

THE ROLE OF INFECTION OF THE LOWER GENITAL TRACT IN THE OUTCOME OF PREGNANCY AND CHILDBIRTH

Tuksanova D.I.,

Bukhara Medical Institute Named after Abu Ali ibn Sino, Bukhara, Uzbekistan.

✓ *Resume*

One of the main directions in solving the problem of safe motherhood is associated with a decrease in reproductive losses, which are considered as the result of the influence of social, medical and biological factors on the health of a pregnant woman, the state of the fetus and newborn.

Key words: infection, genital tract, pregnancy, labor.

Introduktion

The main task of obstetrics at present is to reduce maternal and perinatal morbidity and mortality, in connection with which it is necessary to develop rational programs for the management of patients with various obstetric and perinatal pathologies. One of the main directions in solving the problem of safe motherhood is associated with a decrease in reproductive losses, which are considered as the end result of the influence of social, medical and biological factors on the health of a pregnant woman, the state of the fetus and newborn ().

In recent years, the incidence of inflammatory diseases of the lower part of the urogenital tract has increased, the causative agents of which are found in various associations. According to WHO, more than 300 million new cases are registered in the world every year, including 170 million of trichomoniasis, 89 million of chlamydia.

In our Republic, Uzbekistan also occupies one of the leading places in the structure of obstetric and perinatal morbidity. A significant increase in infections is due to socio - economic changes in the republic, their impact on the social behavior of the population. The increase in the number of genital infections affects the reproductive health of women. Intrauterine infection is one of the most important medical and social problems of modern obstetrics and perinatology. According to many authors,

intrauterine infection develops in 30-50% of children born alive, and in the structure of perinatal losses it accounts for 11-40%.

One of the main directions in solving the problem of safe motherhood is to reduce reproductive losses, which are considered as a result of the unfavorable influence of medical, social and biological factors on the health of a pregnant woman, fetus and newborn.

An especially useful place is occupied by the increase in the frequency of inflammatory diseases of the lower genitals. Most of the pathogens detected are found in associations with which they associate not carrying at any stage of pregnancy, systemic functional disorders of the placenta, intrauterine infection of the fetus, septic complications during childbirth and the postpartum period. This provision was the basis for our study.

Purpose of the study

To study the role of infection of the lower parts of the urogenital tract on the course and outcome of the gestational process.

We examined 50 patients with full-term pregnancies who were admitted for delivery and had clinical manifestations of genital infection (I, main group) and 40 pregnant women with vaginal infection who were admitted for delivery (II, control group).

The age of the surveyed ranged from 18 -430 years. All pregnant women underwent a general clinical, ultrasound examination, as well as a comprehensive microbiological

examination of vaginal discharge, including microscopy of a smear stained according to Gram, and a culture study using conventional methods of isolation and identification of microorganisms. Detection of activation of viral infections -TORCH was carried out by polymerase chain reaction. Sexually transmitted infections (STIs) were excluded in all pregnant women.

The gynecological history was aggravated by chronic adnexitis in 20% of women in the first group and in 10% of the second group; nonspecific vaginitis in 70% and 45%, respectively; erosion of the cervix - in 24% and 15% of pregnant women. The frequency of artificial termination of pregnancy and spontaneous miscarriages in-group I patients was 2.7 times higher than in the control group. Anemia in the main group was detected in 48% of pregnant women, and in the control group - 32% of women.

According to the results of microbiological examination, 32% of pregnant women in the main group were diagnosed with bacterial, 22% - vaginal candidiasis, in 12% of cases there was a combination of bacterial vaginosis and vaginal candidiasis, 18% had nonspecific vaginitis. In all pregnant women of the II group, the state of the vaginal microecosis was regarded as "normoecosis". In-group I, 6% of patients had cytomegalovirus DNA in the scrapings of the cervical canal, and 10% had genital herpes. Ureoplasm was isolated in 28% of patient's in-group I and 10% in-group II.

The course of this pregnancy in every second woman of the main group was complicated by the threat of termination, in 20% of patients - by premature birth, in 14% - by placental insufficiency, in 12% - by polyhydramnios, in 10% of pregnant women with fetal distress, in 6% with fetal development retardation syndrome. These indicators significantly exceeded those in pregnant women in the control group.

Thus, the threat of termination of pregnancy in the control group was noted in 30% of women, premature birth in 12%, placental insufficiency in 6%, polyhydramnios in 8%, fetal distress in 5%, and fetal growth retardation syndrome in 2% of pregnant women.

As for complications in childbirth, in-group I of the examined, premature rupture of amniotic fluid was observed in 24% of pregnant women, in-group II - in 12.5%. Unsatisfactory progress of labor in the main group was found in 14% of women in labor and did not occur in the control group. Emergency operative delivery by cesarean section during labor was performed in 16% of women in labor in the main group and in 7.5% in the control group. Bleeding in the early postpartum period was noted in-group I - in 8% of puerperas, in-group II - in 2.5%. Injuries of the soft birth canal were detected in 48% and 12.5% of cases, respectively.

In the postpartum period, the postpartum women of group I were diagnosed with endometritis (8%), subinvolution of the uterus (10%), wound infection with

the development of infiltrates in the area of sutures on the perineum or their partial divergence (4%). In the control group, there were no cases of endometritis and wound infection. Uterine subinvolution was diagnosed in 2.6% of puerperas. The manifestations of intrauterine infection in the form of rhinitis, conjunctivitis, intrauterine pneumonia, omphalitis were detected in 36% of newborns in-group I and in 15% of newborns in-group II.

Thus, it becomes clear that the problem of genital infection in pregnant women is much more complicated. Than the discomfort associated with the primary episode of the disease, as the risk of serious complications for the mother, fetus and newborn increases significantly. Timely diagnosis during pregnancy, followed by adequate complex therapy, rational management of childbirth, the postpartum period can significantly reduce perinatal morbidity and mortality.

LIST END REFERENCES:

1. Gilbert G.L. Infections in pregnant women. Med J Aust 2002; 176: 229- 236. <https://www.mja.com.au/journal/2002/176/5/1-infections-pregnant-women>;
2. Palasanthiran P, Starr M, Jones C, Giles M; editors. Management of perinatal infections. Sydney: Australasian Society for Infectious Diseases, 2014.
3. Public Health Laboratory Network. Laboratory case definitions for diagnosis of communicable diseases. Canberra: Commonwealth of Australia, 2017.
4. LeBaron C.W., Forghani B, Beck C, et al. Persistence of mumps antibodies after 2 doses of measles?mumps?rubella vaccine. J Infect Dis 2009; 199: 552- 560.
5. Best J.M. Rubella. Semin Fetal Neonatal Med 2007; 12: 182- 192.
6. Craddock?Watson J.E., Ridehalgh MK, Anderson MJ, Pattison JR. Outcome of asymptomatic infection with rubella virus during pregnancy. J Hyg (Lond) 1981; 87: 147- 154.
7. Ahn K.H., Park YJ, Hong SC, et al. Congenital varicella syndrome: a systematic review. J. Obstet. Gynaecol 2016; 36: 563- 566.
8. Gilbert G.L. Parvovirus B19 infection and its significance in pregnancy. Commun Dis. Intel.l 2000; 24 (Suppl): 69- 71.
9. Bascietto F., Liberati M, Murgano D, et al. Outcome of fetuses with congenital parvovirus B19 infection: systematic review and meta?analysis. Ultrasound Obstet Gynecol 2018; 52: 569- 578.
10. Enders M., Klingel K., Weidner A, et al. Risk of fetal hydrops and non?hydropic late intrauterine fetal death after gestational parvovirus B19 infection. J Clin Virol 2010; 49: 163- 171.
11. Verduin E.P., Lindenburg IT, Smits?Wintjens VE, et al. Long?term follow up after intra?uterine transfusions; the LOTUS study. BMC Pregnancy Childbirth 2010; 10: 77- 84.
12. Giorgio E., De Oronzo M.A., Iozza I, et al. Parvovirus B19 during pregnancy: a review. J Prenat Med 2010; 4: 63- 66.
13. Zaripova D.Ya., Sharipova R.G. Efficiency of hormone replacement therapy in prevention menopausal therapy. Academicia an international multidisciplinary research journal. Vol.10, issue 8. Page 422.
14. Zaripova D. and Sharipova R. Comparative evaluation of the use of aleandronic acid in menopausal women complicated by osteoporosis. European journal of biomedical and pharmaceutical sciences. Vol-7, issue-6

Entered 09.11.2020