



New Day in Medicine
Новый День в Медицине

NDM



TIBBIYOTDA YANGI KUN

Ilmiy referativ, marifiy-ma'naviy jurnal



AVICENNA-MED.UZ



ISSN 2181-712X.
EiSSN 2181-2187

4 (66) 2024

Сопредседатели редакционной коллегии:

**Ш. Ж. ТЕШАЕВ,
А. Ш. РЕВИШВИЛИ**

Ред. коллегия:

М.И. АБДУЛЛАЕВ
А.А. АБДУМАЖИДОВ
Р.Б. АБДУЛЛАЕВ
Л.М. АБДУЛЛАЕВА
А.Ш. АБДУМАЖИДОВ
М.А. АБДУЛЛАЕВА
Х.А. АБДУМАЖИДОВ
Б.З. АБДУСАМАТОВ
М.М. АКБАРОВ
Х.А. АКИЛОВ
М.М. АЛИЕВ
С.Ж. АМИНОВ
Ш.Э. АМОНОВ
Ш.М. АХМЕДОВ
Ю.М. АХМЕДОВ
С.М. АХМЕДОВА
Т.А. АСКАРОВ
М.А. АРТИКОВА
Ж.Б. БЕКНАЗАРОВ (главный редактор)
Е.А. БЕРДИЕВ
Б.Т. БУЗРУКОВ
Р.К. ДАДАБАЕВА
М.Н. ДАМИНОВА
К.А. ДЕХКОНОВ
Э.С. ДЖУМАБАЕВ
А.А. ДЖАЛИЛОВ
Н.Н. ЗОЛотова
А.Ш. ИНОЯТОВ
С. ИНДАМИНОВ
А.И. ИСКАНДАРОВ
А.С. ИЛЬЯСОВ
Э.Э. КОБИЛОВ
А.М. МАННАНОВ
Д.М. МУСАЕВА
Т.С. МУСАЕВ
М.Р. МИРЗОЕВА
Ф.Г. НАЗИРОВ
Н.А. НУРАЛИЕВА
Ф.С. ОРИПОВ
Б.Т. РАХИМОВ
Х.А. РАСУЛОВ
Ш.И. РУЗИЕВ
С.А. РУЗИБОВЕВ
С.А.ГАФФОРОВ
С.Т. ШАТМАНОВ (Кыргызстан)
Ж.Б. САТТАРОВ
Б.Б. САФОВЕВ (отв. редактор)
И.А. САТИВАЛДИЕВА
Ш.Т. САЛИМОВ
Д.И. ТУКСАНОВА
М.М. ТАДЖИЕВ
А.Ж. ХАМРАЕВ
Д.А. ХАСАНОВА
А.М. ШАМСИЕВ
А.К. ШАДМАНОВ
Н.Ж. ЭРМАТОВ
Б.Б. ЕРГАШЕВ
Н.Ш. ЕРГАШЕВ
И.Р. ЮЛДАШЕВ
Д.Х. ЮЛДАШЕВА
А.С. ЮСУПОВ
Ш.Ш. ЯРИКУЛОВ
М.Ш. ХАКИМОВ
Д.О. ИВАНОВ (Россия)
К.А. ЕГЕЗАРЯН (Россия)
DONG JINCHENG (Китай)
КУЗАКОВ В.Е. (Россия)
Я. МЕЙЕРНИК (Словакия)
В.А. МИТИШ (Россия)
В.И. ПРИМАКОВ (Беларусь)
О.В. ПЕШИКОВ (Россия)
А.А. ПОТАПОВ (Россия)
А.А. ТЕПЛОВ (Россия)
Т.Ш. ШАРМАНОВ (Казахстан)
А.А. ЩЕГОЛОВ (Россия)
Prof. Dr. KURBANHAN MUSLUMOV (Azerbaijan)
Prof. Dr. DENIZ UYAK (Germany)

**ТИББИЁТДА ЯНГИ КУН
НОВЫЙ ДЕНЬ В МЕДИЦИНЕ
NEW DAY IN MEDICINE**

*Илмий-рефератив, маънавий-маърифий журнал
Научно-реферативный,
духовно-просветительский журнал*

УЧРЕДИТЕЛИ:

**БУХАРСКИЙ ГОСУДАРСТВЕННЫЙ
МЕДИЦИНСКИЙ ИНСТИТУТ
ООО «ТИББИЁТДА ЯНГИ КУН»**

Национальный медицинский
исследовательский центр хирургии имени
А.В. Вишневского является генеральным
научно-практическим
консультантом редакции

Журнал был включен в список журнальных
изданий, рецензируемых Высшей
Аттестационной Комиссией
Республики Узбекистан
(Протокол № 201/03 от 30.12.2013 г.)

РЕДАКЦИОННЫЙ СОВЕТ:

М.М. АБДУРАХМАНОВ (Бухара)
Г.Ж. ЖАРЫЛКАСЫНОВА (Бухара)
А.Ш. ИНОЯТОВ (Ташкент)
Г.А. ИХТИЁРОВА (Бухара)
Ш.И. КАРИМОВ (Ташкент)
У.К. КАЮМОВ (Тошкент)
Ш.И. НАВРУЗОВА (Бухара)
А.А. НОСИРОВ (Ташкент)
А.Р. ОБЛОКУЛОВ (Бухара)
Б.Т. ОДИЛОВА (Ташкент)
Ш.Т. УРАКОВ (Бухара)

4 (66)

2024

апрель

www.bsmi.uz

<https://newdaymedicine.com> E:

ndmuz@mail.ru

Тел: +99890 8061882

Received: 20.03.2024, Accepted: 10.04.2024, Published: 15.04.2024

УДК 616:345-67-234-9/23

FREQUENCY OF THE ZYGOMATIC-ORBITAL FRACTURES AND METHOD OF USING FIBRIN MEMBRANE

Boymuradov Sh.A. <https://orcid.org/2334-4554-8990-0967>

Iminov K.O. <https://orcid.org/6777-6709-8886-0909>

Tashkent Medical Academy 100109, Tashkent, Uzbekistan Farabi Street 2. Tel: +99878 1507825;
E-mail: info@tma.uz

✓ Resume

In several medical fields, there are too many regeneration materials, which prepared from blood one of them is platelet-rich fibrin (PRF)—have been extensively employed due to their significant ability to promote tissue regeneration. Platelets from whole blood were intended to be concentrated using PRP, which was initially proposed a specific restorative technique.

Key words: Platelet-rich fibrin (PRF), Zygomatic-orbital complex, fractures, growth factors, tube, clot, recovery, immune system.

ЧАСТОТА ПЕРЕЛОМОВ СКУЛО-ОРБИТАЛЬНОЙ КОСТИ И МЕТОД ИСПОЛЬЗОВАНИЯ ФИБРИНОВОЙ МЕМБРАНЫ

Боймуратов Ш.А. <https://orcid.org/2334-4554-8990-0967>

Иминов К. О. <https://orcid.org/6777-6709-8886-0909>

Ташкентская Медицинская Академия (ТМА) Узбекистан, 100109, Ташкент, Алмазарский район, ул. Фароби 2, тел: +99878 1507825, E-mail: info@tma.uz

✓ Резюме

В некоторых областях медицины существует слишком много материалов для регенерации, которые готовятся из крови, один из них - богатый тромбоцитами фибрин (PRF) - широко использовался благодаря своей значительной способности способствовать регенерации тканей. Тромбоциты из цельной крови были предназначены для концентрации в PRP, который первоначально был предложен в качестве специфического восстановительного метода.

Ключевые слова: Богатый тромбоцитами фибрин (PRF), скуло-орбитальный комплекс, переломы, факторы роста, трубка, сгусток, восстановление, иммунная система.

ZIGOMATIK ORBITAL SUYAK SINISHI CHASTOTASI VA FIBRIN MEMBRANASIDAN FOYDALANISH USULI

Boymuradov Sh.A. <https://orcid.org/2334-4554-8990-0967>

Iminov K.O. <https://orcid.org/6777-6709-8886-0909>

Toshkent tibbiyot akademiyasi, 100109 Toshkent, O'zbekiston Farobiy ko'chasi 2, Tel: +998781507825 E-mail: info@tma.uz

✓ Rezume

Tibbiyotning ba'zi sohalarida qondan tayyorlangan regeneratsiya materiallari juda ko'p, ulardan biri trombositlarga boy fibrin (PRF) to'qimalarning yangilanishini rag'batlantirish qobiliyati tufayli keng qo'llanilgan. To'liq qon trombositlari dastlab o'ziga xos reduktiv usul sifatida taklif qilingan PRP konsentratsiya qilish uchun mo'ljallangan.

Kalit so'zlar: fibrin trombositlariga boy (PRF), zigomatik orbital kompleks, yoriqlar, o'sish omillari, naycha, pihti, tiklanish, immunitet tizimi.

Introduction

Vascular endothelial growth factor (VEGF), transforming growth factor beta (TGF- β), Platelet-derived growth factor (PDGF) and are just a few of the naturally produced autologous growth factors found in PRP. This group of growth factors is in charge of encouraging the angiogenesis (the development of new blood vessels) as well as the migration and proliferation of different cell types [1]. However, utilizing a centrifuge to sort out blood-derived growth factors and cells. after peripheral blood collection is straightforward and inexpensive, and it has long been recognized as an efficient and convenient way to extract natural growth influencing variables tissue regeneration. In order to remove anticoagulant, PRF was subsequently created as a second-generation platelet concentrate [2]. Because leukocytes are wanted to be included, this concentration is sometimes. Identified as leukocytes and platelet-rich fibrin (L-PRF). This is because a 700g for 12 min procedure is typically used [3-4]. The removal of anticoagulants causes blood in the blood collection tube to clot over time. As a result, centrifugation must be started as soon as possible by the treating physician in order to separate the blood layers. One significant benefit is fibrin plates generated in platelet rich fibrin, In contrast to PRP, this encourages the gradual and constant release of growth factors throughout time. [8]. Additionally, a liquid-PRF (injectable-PRF, or i-PRF) with a higher concentration of platelets and leukocytes was created by lowering centrifugation speeds and times [5]. It was recently revealed that, when compared to the outcomes of standard fixed-angle centrifuges used to create solid-PRF, the horizontal centrifugation of PRF was superior in accumulating platelets and leukocytes. Platelet/leukocyte counts and/or concentrations were improved by up to 3.

Purpose of the study: To investigate the incidence of fractures of the skeletal-orbital bone and the method of using the fibrine membrane

Methods and materials

Before collecting blood samples, 29 patients supplied prior permission. All methods involving human subjects in this study adhered to the ethical standards set by the organizational as well as national scientific committees. All blood samples were taken at the Tashkent Medical Academy and used in compliance with Medical Ethics Standards and Guidelines. Factors influencing Genetics, acquired factors (e.g., platelet activation, hyperhomocysteinemia, abnormal thrombin and factor XIII levels in plasma, blood flow, oxidative stress, hyperglycemia, medications, and cigarette smoking), and environmental factors (e.g., temperature, reducing agents, chloride and calcium ion concentrations) all contribute to fibrin clot formation and structure. All patients confirmed that they did not have any of the aforementioned conditions. Prior to the studies, the CBCs were analyzed to confirm conventional cell count ranges. The PRF was produced using an Dilab centrifuge. Each of the 29 participants supplied 29 vials of blood in normal 10 mL glass collection tubes (vacutainers) [7]. We lead to common blood test till preparing PRF membranes. Unique side of this method is we complete all steps PRF preparation. Moreover, those steps we do in operation theatre simultaneously with performing reconstruction zygomatic orbital complex. During this performance surgeon use titanium plates for reconstruction the floor of orbital after “blow-out” fractures [8-12]. Titanium plate covered with fibrin membranes and cytokines produce from PRF then they useful for recovery bone structures of ZOC and regeneration soft tissue.

Result and discussions

As in the preceding group, all (n = 29) patients with varied SOC fractures who were treated with autologous fibrin membranes supplemented with platelets were classified strictly using the Manson 1990 classification.




Figure 1 Distribution of patients with ZOC fractures. Treatment of ZOC with the use of autologous fibrin membrane enriched with platelets according to the Manson classification.

Maxillofacial surgeons, ophthalmologists, and, if necessary, neurosurgeons examined the patients' conditions prior to surgery. In the second group (n = 29), all reconstructive procedures on the zygomatic-orbital complex were carried out making use of PRF membranes.




Diagram 1 complaints of patients before surgery on the zygomatic-orbital complex.

The diagram shows that in the main group of the study, patients before the surgical period complained more than 25 about the aesthetic defect of the zygomatic-orbital complex, 22 complaints were related to enophthalmos, hypophthalmos was identified in 19 patients, the step symptom was identified in 15 patients, it is important to note that all complaints have a combined character. One of the features of our work, in addition to the general standard laboratory tests that the patient underwent in the emergency room, before each preparation of PRF membranes, we conducted a thorough general blood test with a detailed formula of 31 indicators on a hematological analyzer manufactured in 2022 by DF 50 DYMIN, which complies with the European standard in operating rooms. After an examination, the maxillofacial surgeon at the Tashkent Medical Academy's plastic surgery department orders laboratory and instrumental studies for patients, such as computed tomography with a 3D model of the skull (to determine the nature of the fracture, the volume of surgical manipulations which performed, and the size of the lower wall of the orbit). Laboratory tests include a general blood test (to determine the number of platelets, as it is difficult to obtain a high-quality fibrin membrane from a small number of platelets) and a coagulogram (particularly to determine the amount of fibrinogen, INR, and APTT), as well as standard studies such as blood biochemistry and blood tests for infection. An anesthesiologist is examined after obtaining anthropometric indicators by analyzing photographs of patients and MSCT data to determine the level of enophthalmos of the affected eye socket (enophthalmos is an ophthalmological pathology in which the eyeball sinks excessively into the socket in which the orbit is present). All SVC restoration procedures involve general anesthesia and tracheal intubation. The temperature in the operating room is regulated to 21-22°C once the patient is intubated (since ambient temperature is the most important component for separating fibrin membranes). The surgical field is processed, the surgical incision is made with a disposable surgery scalpel, exposing the fracture lines of the zygomatic bone along the zygomaticofrontal suture and the lower orbital edge through the skin incision, retreating 5 mm from the edge of the lower eyelash line, then coagulation of the vessels is performed, fixation of the upper edge of the surgical wound with a ligature, and this causes the muscle layer of the extraocular muscles to After reaching the bone structure of the eye socket, the eyeball is gently elevated using a short medical equipment to maximize visibility. Before administering anesthesia, venous blood is drawn from the ulnar region. After placing the tubes without anticoagulants at an angle of 45°, immediately start the centrifugation stage with a volume of 9-10 ml. The centrifugation step lasts 8 minutes and takes place at 1300 rpm. After centrifugation, the test tube is placed on a stand, and the room temperature should be 21-22°C to allow the fibrin plate to come into contact with oxygen in the air. The upper part of the

resultant material, which is seldom plasma, must be removed using a syringe. Contact with oxygen must last at least 6 minutes. Using medical devices (tweezers), the fibrin membrane (PRF) with the red blood cell mass is gently pulled from the test tube, and the PRF is precisely separated from the red blood cell mass. Following separation, a liquid component may be visible in the fibrin membrane, which includes a considerable number of leukocytes. To achieve a pure fibrin membrane, PRF is compressed with a metal plate for 10-15 minutes.

Conclusion

Analysis the results of traditional methods treatment of injuries the zygomatic orbital complex shown a high level of postoperative complications (20%) such as paraorbital edema, a feeling of discomfort when moving the eyeball, and severe symptoms of inflammation, caused by insufficient administration of anti-inflammatory drugs in the postoperative period and features of the course diseases due to the development of dysfunction in the zygoma. The use of PRF membranes with titanium polymer compounds in ZOC repair has resulted in improved postoperative outcomes.

According to the literature research, no method for preparing a PRF membrane intraoperatively in reconstructive procedures on the zygomatic-orbital complex has been created.

The widespread use of PRF membrane intraoperatively in reconstructive procedures on the zygomatic-orbital complex reduces the percentage of postoperative complications and the proportion of radical operations on the ZOC, resulting in a 1.5-fold reduction in inpatient patient days in hospital.

LIST OF REFERENCES:

1. Delaini F, Poggi A, Donati MB. Enhanced affinity for arachidonic acid in platelet-rich plasma from rats with Adriamycin-induced nephrotic syndrome. // *Thromb Haemost.* 1982;48(3):260–2.
2. Miron RJ, Zucchelli G, Pikos MA, Salama M, Lee S, Guillemette V, Fujioka, Kobayashi M, Bishara M, Zhang Y, Wang H-L, Coi J. Use of platelet-rich fibrin in regenerative dentistry: a systematic review. // *Clin Oral Invest.* 2017;21(6):1913–27.
3. Miron RJ, Choukroun J, Ghanaati S. Necessity for standardization of relative centrifugal force values in studies on platelet rich fibrin; response to letter to the editor. // *J Periodontol.* 2018;90:122–5.
4. Ehrenfest DMD, Rasmusson L, Albrektsson TJ. Tib: classification of platelet concentrates: from pure platelet-rich plasma (P-PRP) to leucocyte-and platelet-rich fibrin (L-PRF). // *Trends Biotechnol.* 2009;27(3):158–67.
5. Kobayashi E, Fluckiger L, Fujioka-Kobayashi M, Sawada K, Sculean A, Schaller B, Miron RJ. Comparative release of growth factors from PRP, PRF, and advanced-PRF. // *Clin Oral Invest.* 2016;20(9):2353–60.
6. Miron RJ, Fujioka-Kobayashi M, Hernandez M, Kandalam U, Zhang Y, Ghanaati S, Choukroun J. Injectable platelet rich fibrin (i-PRF): opportunities in regenerative dentistry? // *Clin Oral Invest.* 2017;21:2619–27.
7. Miron RJ, Chai J, Zheng S, Feng M, Sculean A, Zhang Y. A novel method for evaluating and quantifying cell types in platelet rich fibrin and an introduction to horizontal centrifugation. // *J Biomed Mater Res, Part A.* 2019;107(10):2257–71.
8. Nunes CR, Roedersheimer M, Simske S, Luttgies M. Effect of microgravity, temperature, and concentration on fibrin and collagen assembly. // *Microgravity Sci Technol.* 1995;8(2):125–30.
9. Ghanaati S, Booms P, Orlowska A, Kubesch A, Lorenz J, Rutkowski J, Landes C, Sader R, Kirkpatrick C, Choukroun J. Advanced platelet-rich fibrin: a new concept for cell-based tissue engineering by means of inflammatory cells. // *J Oral Implantol.* 2014;40(6):679–89.
10. Takahashi A, Tsujino T, Yamaguchi S, Isobe K, Watanabe T, Kitamura Y, Okuda K, Nakata K, Kawase T: Distribution of platelets, transforming growth factor-beta1, platelet-derived growth factor-BB, vascular endothelial growth factor and matrix metalloprotease-9 in advanced platelet-rich fibrin and concentrated growth factor matrices. // *J Investig Clin Dent* 2019:e12458.
11. Miron RJ, Dham A, Dham U, Zhang Y, Pikos MA, Sculean A. The effect of age, gender, and time between blood draw and start of centrifugation on the size outcomes of platelet-rich fibrin (PRF) membranes. // *Clin Oral Invest.* 2018;23:2179–85.
12. Varela HA, Souza JCM, Nascimento RM, Araujo RF Jr, Vasconcelos RC, Cavalcante RS, Guedes PM, Araujo AA. Injectable platelet rich fibrin: cell content, morphological, and protein characterization. // *Clin Oral Invest.* 2019;23(3):1309–18

Entered 20.03.2024

