




New Day in Medicine
Новый День в Медицине

NDM



TIBBIYOTDA YANGI KUN

Ilmiy referativ, marifiy-ma'naviy jurnal



AVICENNA-MED.UZ



ISSN 2181-712X.
EiSSN 2181-2187

9 (71) 2024

**Сопредседатели редакционной
коллегии:**

**Ш. Ж. ТЕШАЕВ,
А. Ш. РЕВИШВИЛИ**

Ред. коллегия:

М.И. АБДУЛЛАЕВ
А.А. АБДУМАЖИДОВ
Р.Б. АБДУЛЛАЕВ
Л.М. АБДУЛЛАЕВА
А.Ш. АБДУМАЖИДОВ
М.А. АБДУЛЛАЕВА
Х.А. АБДУМАЖИДОВ
Б.З. АБДУСАМАТОВ
М.М. АКБАРОВ
Х.А. АКИЛОВ
М.М. АЛИЕВ
С.Ж. АМИНОВ
Ш.Э. АМОНОВ
Ш.М. АХМЕДОВ
Ю.М. АХМЕДОВ
С.М. АХМЕДОВА
Т.А. АСКАРОВ
М.А. АРТИКОВА
Ж.Б. БЕКНАЗАРОВ (главный редактор)
Е.А. БЕРДИЕВ
Б.Т. БУЗРУКОВ
Р.К. ДАДАБАЕВА
М.Н. ДАМИНОВА
К.А. ДЕХКОНОВ
Э.С. ДЖУМАБАЕВ
А.А. ДЖАЛИЛОВ
Н.Н. ЗОЛотова
А.Ш. ИНОЯТОВ
С. ИНДАМИНОВ
А.И. ИСКАНДАРОВ
А.С. ИЛЬЯСОВ
Э.Э. КОБИЛОВ
А.М. МАННАНОВ
Д.М. МУСАЕВА
Т.С. МУСАЕВ
М.Р. МИРЗОЕВА
Ф.Г. НАЗИРОВ
Н.А. НУРАЛИЕВА
Ф.С. ОРИПОВ
Б.Т. РАХИМОВ
Х.А. РАСУЛОВ
Ш.И. РУЗИЕВ
С.А. РУЗИБОВЕВ
С.А.ГАФФОРОВ
С.Т. ШАТМАНОВ (Кыргызстан)
Ж.Б. САТТАРОВ
Б.Б. САФОВЕВ (отв. редактор)
И.А. САТИВАЛДИЕВА
Ш.Т. САЛИМОВ
Д.И. ТУКСАНОВА
М.М. ТАДЖИЕВ
А.Ж. ХАМРАЕВ
Д.А. ХАСАНОВА
А.М. ШАМСИЕВ
А.К. ШАДМАНОВ
Н.Ж. ЭРМАТОВ
Б.Б. ЕРГАШЕВ
Н.Ш. ЕРГАШЕВ
И.Р. ЮЛДАШЕВ
Д.Х. ЮЛДАШЕВА
А.С. ЮСУПОВ
Ш.Ш. ЯРИКУЛОВ
М.Ш. ХАКИМОВ
Д.О. ИВАНОВ (Россия)
К.А. ЕГЕЗАРЯН (Россия)
DONG JINCHENG (Китай)
КУЗАКОВ В.Е. (Россия)
Я. МЕЙЕРНИК (Словакия)
В.А. МИТИШ (Россия)
В.И. ПРИМАКОВ (Беларусь)
О.В. ПЕШИКОВ (Россия)
А.А. ПОТАПОВ (Россия)
А.А. ТЕПЛОВ (Россия)
Т.Ш. ШАРМАНОВ (Казахстан)
А.А. ЩЕГОЛОВ (Россия)
Prof. Dr. KURBANHAN MUSLUMOV (Azerbaijan)
Prof. Dr. DENIZ UYAK (Germany)

**ТИББИЁТДА ЯНГИ КУН
НОВЫЙ ДЕНЬ В МЕДИЦИНЕ
NEW DAY IN MEDICINE**

*Илмий-рефератив, маънавий-маърифий журнал
Научно-реферативный,
духовно-просветительский журнал*

УЧРЕДИТЕЛИ:

**БУХАРСКИЙ ГОСУДАРСТВЕННЫЙ
МЕДИЦИНСКИЙ ИНСТИТУТ
ООО «ТИББИЁТДА ЯНГИ КУН»**

Национальный медицинский
исследовательский центр хирургии имени
А.В. Вишневского является генеральным
научно-практическим
консультантом редакции

Журнал был включен в список журнальных
изданий, рецензируемых Высшей
Аттестационной Комиссией
Республики Узбекистан
(Протокол № 201/03 от 30.12.2013 г.)

РЕДАКЦИОННЫЙ СОВЕТ:

М.М. АБДУРАХМАНОВ (Бухара)
Г.Ж. ЖАРЫЛКАСЫНОВА (Бухара)
А.Ш. ИНОЯТОВ (Ташкент)
Г.А. ИХТИЁРОВА (Бухара)
Ш.И. КАРИМОВ (Ташкент)
У.К. КАЮМОВ (Тошкент)
Ш.И. НАВРУЗОВА (Бухара)
А.А. НОСИРОВ (Ташкент)
А.Р. ОБЛОКУЛОВ (Бухара)
Б.Т. ОДИЛОВА (Ташкент)
Ш.Т. УРАКОВ (Бухара)

9 (71)

2024

сентябрь

www.bsmi.uz

https://newdaymedicine.com E:

ndmuz@mail.ru

Тел: +99890 8061882

Received: 20.08.2024, Accepted: 02.09.2024, Published: 10.09.2024

UDK 616.98-036-07-08:578.834.11

IMPROVEMENT OF PRIMARY PROPHYLAXIS AND TREATMENT OF SPONTANEOUS BACTERIAL PERITONITIS COMPLICATED IN VIRUS ETIOLOGY LIVER CIRRHOSIS

Oblokulov A.R. <https://orcid.org/0000-0002-6891-359X>
Mukhammadieva M.I. <https://orcid.org/0009-0004-6329-4883>

Bukhara State Medical Institute named after Abu Ali ibn Sino, Uzbekistan, Bukhara c.
Gijduvan street 23, e-mail: info@bsmi.uz

✓ Resume

The article presents clinical and laboratory data of the examined patients. 120 patients were under observation. Main group I included 60 patients suffering from liver cirrhosis of viral etiology with SBP, main group II included 60 patients with liver cirrhosis. As a control, 20 practically healthy people were also examined.

In all patients with SBP (Group 1) at the stage of decompensation of liver cirrhosis of viral etiology, the level of PCT was significantly higher by 10 times, amounting to 0.88 ± 0.04 , compared with patients with uncomplicated SBP (Group 2), in which is equal to 0.08 ± 0.02 ($p=0.05$). It was found that the level of CRP in the 1st group was 32.4 ± 8.23 and increased by 3.75 times ($p=0.05$) compared with the 2nd group.

Based on these values, it can be concluded that the level of PCT and CRP in the blood serum has been proposed as a marker for early non-invasive diagnosis in patients with cirrhosis and SBP.

Key words: liver cirrhosis, spontaneous bacterial peritonitis, procalcitonin, C-reactive protein.

СОВЕРШЕНСТВОВАНИЕ ПЕРВИЧНОЙ ПРОФИЛАКТИКИ И ЛЕЧЕНИЯ СПОНТАННОГО БАКТЕРИАЛЬНОГО ПЕРИТОНИТА, ОСЛОЖНЕННОГО ЦИРРОЗОМ ПЕЧЕНИ ВИРУСНОЙ ЭТИОЛОГИИ

Облокулов А.Р. <https://orcid.org/0000-0002-6891-359X>
Мухаммадиева М.И. <https://orcid.org/0009-0004-6329-4883>

Бухарский государственный медицинский институт им. Абу Али ибн Сино, Узбекистан, г. Бухара, улица Гиждуванская, 23, e-mail.ru: info@bsmi.uz

✓ Резюме

В статье представлены клиничко-лабораторные данные обследованных пациентов. Под наблюдением находились 120 пациентов. В I основную группу вошли 60 больных циррозом печени вирусной этиологии с СБП, во II основную группу — 60 больных циррозом печени. В качестве контроля также были обследованы 20 практически здоровых людей.

У всех больных СБП (I-я группа) на стадии декомпенсации цирроза печени вирусной этиологии уровень ПКТ был достоверно выше в 10 раз, составив $0,88 \pm 0,04$, по сравнению с больными неосложненным СБП (2-я группа), в что равно $0,08 \pm 0,02$ ($p=0,05$). Установлено, что уровень СРБ в I-й группе составил $32,4 \pm 8,23$ и увеличился в 3,75 раза ($p=0,05$) по сравнению со 2-й группой.

На основании этих значений можно сделать вывод, что уровень ПКТ и СРБ в сыворотке крови предложен в качестве маркера ранней неинвазивной диагностики у больных циррозом печени и СБП.

Ключевые слова: цирроз печени, спонтанный бактериальный перитонит, прокальцитонин, С-реактивный белок.

VIRUS ETIOLOGIYALI JIGAR SIRROZIDA ASORATLANGAN SPONTAN BAKTERIAL PERITONITNI BIRLAMCHI PROFILAKTIKASI VA DAVOSINI TAKOMILLASHTIRISH

Obloqulov A.R. <https://orcid.org/0000-0002-6891-359X>
Muxammadiyeva M.I. <https://orcid.org/0009-0004-6329-4883>

Abu Ali ibn Sino nomidagi Buxoro davlat tibbiyot instituti, Ozbekiston, Buxoro sh.
Gijduvon kochasi, 23 e-mail: info@bsmi.uz

✓ Rezyume

Maqolada tekshirilgan bemorlarning klinik va laboratoriya ma'lumotlari keltirilgan. 120 nafar bemor kuzatuv ostida edi. I asosiy guruhga SBP bilan virusli etiologiyali jigar sirrozi bilan og'rigan 60 nafar bemor, II asosiy guruhga jigar sirrozi bilan og'rigan 60 nafar bemor kiritilgan. Nazorat tariqasida 20 nafar amalda sog'lom odam ham tekshirildi.

Virusli etiologiyali jigar sirrozi dekompensatsiyasi bosqichida SBP (1-guruh) bo'lgan barcha bemorlarda PKT darajasi asoratlanmagan SBP (2-guruh) bo'lgan bemorlarga nisbatan 10 baravar yuqori bo'lib, $0,88 \pm 0,04$ ni tashkil etdi. $0,08 \pm 0,02$ ga teng ($p=0,05$). 1-guruhdagi CRO darajasi $32,4 \pm 8,23$ ni tashkil etgani va 2-guruhga nisbatan 3,75 martaba ($p=0,05$) oshganligi aniqlandi.

Ushbu qiymatlarga asoslanib, qon zardobidagi PKT va CRO darajasi siroz va SBP bilan og'rigan bemorlarda erta invaziv bo'lmagan diagnostika uchun marker sifatida taklif qilingan degan xulosaga kelish mumkin.

Kalit so'zlar: jigar sirrozi, spontan bakterial peritonit, prokalsitonin, C-reaktiv oqsil.

Relevance

The World Health Organization (WHO) estimates that more than 325 million people worldwide are infected with viral hepatitis B or C and 1.4 million people die each year. In terms of mortality, hepatitis B and C together are second only to tuberculosis, and the number of people infected with hepatitis is 9 times higher than the number of people infected with HIV. Yet over 80% of people with hepatitis do not have access to prevention, testing and treatment [1,2].

Despite the successes achieved in the fight against many infectious diseases, in modern medicine the problem of chronic viral hepatitis (CVH) in Uzbekistan, as well as throughout the world, continues to be relevant [3,4]. The significance of this problem is determined not only by the ubiquitous and significant distribution, but also by the predominant lesion of people of working age, a long and progressive course, the complexity of therapy and patient management, severe consequences, such as cirrhosis of the liver (LC) and hepatocellular carcinoma (HCC), as well as significant costs of medical and social support for this category of patients [5, 6].

Spontaneous bacterial peritonitis is a common and severe complication. About 32-40% of hospitalized patients with cirrhosis develop bacterial infections at presentation or during hospitalization. The problems of their diagnostics are waiting for their solution, which have not been solved so far [7,8].

1.A bacterial infection is either present on admission or develops during hospitalization in approximately 30% of patients with cirrhosis [9], and the most common form of these infections is spontaneous bacterial peritonitis (SBP) [10,11].

Patients with cirrhosis have altered protection against bacteria associated with a decrease in bacterial clearance [12]. This immune defect promotes bacterial translocation caused by increased intestinal permeability and bacterial overgrowth [13]. Diagnosis of bacterial complications of cirrhosis is often difficult due to the erased clinical picture of the disease. Sometimes infectious complications appear only as an aggravation of hepatic encephalopathy. Simple and affordable screening tests for bacterial infection in liver cirrhosis include C-reactive protein (CRP) and procalcitonin [14].

Procalcitonin has been proposed in studies as a potentially valuable serum biomarker for diagnosing bacterial infections in general and SBP in particular [15].

CRP is a prominent biochemical marker of inflammation caused by a variety of causes, including infectious and non-infectious inflammatory diseases, and has also been shown to be involved in several immunological functions [16,17].

Thus, infections are common in patients with cirrhosis of the liver, and SBP is one of the most common, with variable frequency but significant mortality. One of the most important factors in the management of

this significant consequence of decompensated liver cirrhosis is early detection. It is critical to find non-invasive, accessible, and easy-to-apply SBP-related parameters that play a predictive role. However, it should be borne in mind that these methods cannot completely replace paracentesis; More research is needed to determine if non-invasive methods are accurate enough to detect the development of SBP in cirrhosis.

Purpose of the study: To develop additional studies to determine whether non-invasive methods are sufficiently accurate to detect the development of SBP in cirrhosis.

Materials and methods

Clinical observations, laboratory and instrumental examinations of patients with liver cirrhosis of viral etiology were carried out in the Bukhara Regional Infectious Diseases Hospital, at the Research Institute of Virology of the Republican Specialized Scientific and Practical Medical Center for Epidemiology, Microbiology, Infectious and Parasitic Diseases.

Main group I included 60 patients suffering from liver cirrhosis of viral etiology with SBP, main group II included 58 patients with liver cirrhosis. As a control, 20 practically healthy people were also examined.

The diagnosis of cirrhosis with viral etiology was established on the basis of the epidemiological anamnesis, medical history, clinical data and on the basis of laboratory data.

In order to establish the viral etiology of the diagnosis of cirrhosis, an analysis was made to determine the markers of infection with the HBV, HDV viruses, HCV was determined by the polymerase chain reaction (PCR).

Among them, in the 1st group, HBV- infection was observed in 15 (25%) patients, HCV infection - in 25 (41.7%), HBV + HCV- infection - in 1 (1.67%), HBV + HDV- infection - in 19 (31.7%). In the second group, HBV infection was noted in 13 (22.4%) patients, HCV infection in 27 (46.6%) patients, HBV+HCV- infection in 1 (1.72%), HBV+HDV- infection in 17 (29.3%)

Result and discussion

The results of the study showed that in the first group of 60 patients, 38 (63.3%) were male and 22 (36.7%) female, and in the second of 58 patients, 27 (46.55%) were male and 31 (53.4%) consisted of women. It should be noted that the probability of occurrence of SBP prevailed in men.

The average age index of sick men of the 1st group was 48.4±10.1 years, and in women 43.1±14.8 years. The age indicators of patients of the 2nd group did not differ from the first and amounted to 45.5±10.1 and 52.1±13.07, respectively.

The duration of the disease in patients of both groups ranged from one month to several years, while the majority of them, 46.7% and 41.4%, respectively, indicated that liver cirrhosis was diagnosed several years ago.

An analysis of the incidence of the disease between groups showed that in group 1, cirrhosis was most often detected in 17 (28%) people aged 51-60 years, the bulk of which were 12 (20%) men and 5 (8.3%) women. Unlike the first group, in the 2nd group cirrhosis of the liver was detected in the age group of 41-50 years, mainly in 10 (17%) men and 6 (10%) women.

These data correspond to the results of foreign epidemiological studies, in particular, the results of V.T. Ivashkin, which included more than 50,000 patients from different regions of the Russian Federation (2014) [18].

The main clinical signs of the examined patients were general weakness in 102 (86.4%), decreased appetite in 74 (62.7%), nausea and vomiting in 51 (43.2%), pruritus in 25 (21.2%) and abdominal pain 71 (60.2%), 70 (59.3%) had subfebrile fever, 19 (16.1%) had a high temperature, 50 (42.4%) had severe jaundice, 43 (36.4%) - moderate jaundice, 71 (60.2%) - splenomegaly, 61 (51.7%) were found to have edema (Picture 2). These data are consistent with the results of foreign clinical trials [19,20].

When examining patients, "spider veins" were noted in 75 (63.6%), palmar erythema in 43 (36.4%), varicose veins of the esophagus in 68 (57.6%). Cases of epistaxis were observed in 47 (39.8%) patients.

In addition, a comparative analysis was made of the duration of complaints and symptoms in patients in the comparison groups, which revealed significant differences between them. The results of a comparative analysis showed that patients of the 2nd group had more poor symptoms, in contrast to patients with SBP (1st group). The leading symptoms in patients of this group were itching (52±15.6%) and splenomegaly according to liver ultrasound, mostly asymptomatic (Table 1).

The majority of patients in the second group had asthenovegetative syndrome, which was characterized by the presence of general weakness and decreased performance (84.1 ± 9.03% of the total number of the

group), while in patients of the first group, these signs occurred with a frequency 89.4±8.3%. In 72.3±15.4% of patients of the second group, dyspeptic symptoms were noted, in particular, episodic occurrence of heartburn, heaviness in the epigastric region, decreased appetite and loosening of the stool. Pain in the abdomen in patients of the 2nd group was 65±14%, and in patients of the 1st group, this figure was 76±13%.

Tab 1.
Average duration of subjective complaints of patients in comparison groups (M±m)

Groups	Malaise	Abdominal pain	Nausea, vomiting	Loss of appetite	Skin itching
1st group (n=60)	8,4±2,7	7,2±2,7	6,5±2,4	7,19±2,76	5,48±3,23
2nd group (n=58)	8,3±1,9	6,5±2	7,2±2,2	7,12±1,96	5,12±2,09

An objective examination was organized to identify signs such as jaundice of the skin and sclera of the eyes, white coating on the surface of the tongue, the presence of “caput medusae” and liver symptoms such as palmar erythema, spider veins, bleeding from the gums and nose.

In case of liver diseases, the skin acquires a yellow color of varying intensity: from dark gray to dark yellow, which in turn occurs as a result of the deposition of bilirubin and its metabolic products in the skin. Jaundice often affects the eyes and mucous membranes: instead of white, they turn yellow.

An objective examination of the patients revealed that in the 1st group during the treatment period, icterus of the skin and sclera of the eyes lasted 8.35±2.17 and 9.92±3.16 days, respectively, while the duration of these complaints in patients was 2- th group was 6.33±2.69 and 6.11±2.28, respectively, which was significantly less than the first group.

Palmar erythema or "hepatic" palms is a condition in which there is even redness of both palms. Palmar erythema appears due to circulatory disorders in the portal vein, leading to the formation of arteriovenous anastomoses. Reddening of the palms occurs when additional vascular branches are formed that connect the veins and arteries for the normal nutrition of the skin and other organs.

Signs of palmar erythema in patients of the 1st group lasted on average 1 day longer compared to the 2nd group, amounting to 9.63±2.68 and 8.93±1.98 days, respectively. The difference in spider vein symptom scores had similar results, with the duration being 9.28±0.68 and 8.13±2.16 days. Also, the appearance of spider veins in patients with cirrhosis may be associated with a violation of the estrogen conformational properties of the cytochrome system in the damaged liver.

Patients in whom cirrhosis proceeded with the development of complications in the form of the development of SBP (Group 1), in contrast to patients in whom cirrhosis of the liver proceeded without the development of complications in the form of SBP (Group 2), hypoalbuminemia was observed, the concentration of total protein in the blood in of the first group of patients was in the range of 27.2±2.3, while in patients of the second group this indicator was 31.2±4.7. This indicated a decrease in the protein-forming function in patients of both groups.

In addition, the mean values of the amount of fibrinogen in patients with SBP (Group 1) 0.73±0.14 were significantly lower by 3.83 times than in the group of patients without SBP (Group 2) 2.8±0.8 (p<0.05). It should be noted that the level of fibrinogen in the 2nd group fluctuated within the minimum values. Hypofibrinogenemia develops due to a decrease in the synthesis of fibrinogen, its increased consumption, as well as increased destruction during the activation of the fibrinolysis process.

Diagnosis of bacterial complications of cirrhosis is often difficult due to the blurred clinical picture of the disease. Sometimes infectious complications appear only as an aggravation of hepatic encephalopathy. Simple and affordable screening tests for bacterial infection in cirrhosis include C-reactive protein and procalcitonin.

In all patients with SBP (Group 1) at the stage of decompensation of liver cirrhosis of viral etiology, the level of PCT was significantly higher by 10 times, amounting to 0.88±0.04, compared with patients with uncomplicated SBP (Group 2), in which is equal to 0.08±0.02 (p=0.05).

When analyzing the increase in the level of PCT in the blood serum in patients of the first group, the content of PCT was observed in the range of 0.2-0.3 ng/ml in 26.7% (16 people), in the range of 0.2-0.3 ng/ml in 36.8% (21 people) 0.3-0.5 ng/ml, 21.7% (13 people) 0.5-1.0 ng/ml, and 14.8% (10 people) 1.0 ng/ml ml, in patients of the control group it was registered in 74.2% (43 people) 0.05-0.1 ng/ml, in 25.8%

(15 people) it was 0.1-0.2 ng/ml (Fig. 2). In 2013 Su DH et al. performed a meta-analysis of PCT including 181 episodes with suspected infection, resulting in pooled sensitivity, specificity, positive likelihood ratios, and negative likelihood ratios of 86%, 80%, 7.73, and 0.14, respectively, suggesting PCT as a diagnostic aid for SPB with moderate to high accuracy [21]. In another 2015 study, Yang Y.'s meta-analysis included 742 patients across 17 studies and found that PCT at a cutoff of 0.42–0.76 ng/mL has a sensitivity of 82% and a specificity of 86% for predicting SBP [22].

According to the results of the analysis, an increase in the PCT inflammatory marker of 0.2 ng/ml in patients complicated by SBP indicates the addition of a bacterial infection and is an indication for prescribing antibacterial drugs.

As can be seen from the data presented, there were statistically significant differences in the values of laboratory parameters characterizing the functional state of the liver between patients of groups I and II.

The final stage of biochemical analysis in the studied blood groups was the study of the amount of C-reactive protein. It was found that the level of CRP in the 1st group was 32.4 ± 8.23 and increased by 3.75 times ($p=0.05$) compared with the 2nd group. An increase in the level of C-reactive protein in the blood indicated significant tissue damage, inflammation, infection, and the presence of a virus. According to several studies, the threshold level of CRP for the diagnosis of infection in patients with liver cirrhosis should be between 2.0 and 8.0 mg/dL [23, 24].

Conclusions

Based on these values, it can be concluded that the level of PCT and CRP in the blood serum has been proposed as a marker for early non-invasive diagnosis in patients with cirrhosis and SBP.

LIST OF REFERENCES:

1. Conn H.O. Spontaneous peritonitis and bacteremia in Laennec's cirrhosis caused by enteric organisms. A relatively common but rarely recognized syndrome // *Ann. Intern. Med.* 1964;60:568-5802.
2. Винницкая Е.В., Осипов Г.А., Дроздов В.Н. и др. Диагностика спонтанного бактериального перитонита при циррозе печени // *Экспериментальная и клиническая гастроэнтерология.* 2008;3:18-24.
3. Jepsen P. et al. Prognosis of patients with liver cirrhosis and spontaneous bacterial peritonitis // *Hepatogastroenterology.* 2003;50:2133-2136.
4. Merino Rodriguez B. et al. Prognostic value of a spontaneous bacterial peritonitis episode in cirrhotic patients in clinical practice // *Gastroenterol. Hepatol.* 2008;31(9):560-565.
5. Бондаренко В.М., Воробьев А.А. Дисбиозы и препараты с пробиотической функцией // *Журн. Микробиол.* 2004;1:84-92.
6. Ghassemi S., Garsia-Tsao G. Prevention and treatment of infections in patients with cirrhosis // *Best Pract. Res. Clin. Gastroenterol.* 2007;21(1):77-93.
7. Облокулов А.Р., Нарзиев И.И., Абдуллаев Т.У., Облокулов А.А. ВГВ лямблиоз билан кечган микс инфекцияда алергик кўрсаткичларнинг аҳамияти // *Тиббиётда янги кун.* 2018;2(22):101-104.
8. Белобордова Н.В., Осипов Г.А. Гомеостаз малых молекул микробного происхождения и его роль взаимоотношениях микроорганизмов с хозяином // *Вестник РАМН.* 1999;16(7):25-31.
9. Ong H.S., Kang C.I., Lee J.A. et al. Clinical significance and outcome of nosocomial acquisition of spontaneous bacterial peritonitis in patients with liver cirrhosis. // *Clin Infect Dis.* 2009;48(9):1230-1236.
10. D'Amico G., Garsia-Tsao G., Pagliaro L. Natural history and prognostic indicators of survival in cirrhosis: a systematic review of 118 studies. // *J Hepatol.* 2006;44(1):217-231.
11. Mukhammadieva M.I. (2022). Modern Clinical and Biochemical Characteristics of Liver Cirrhosis Patients of Viral Etiology with Spontaneous Bacterial Peritonitis // *Texas Journal of Medical Science.* 2022; P.86-90.
12. Mukhammadieva M.I. (2023). Вирус этиологияли жигар циррози беморларида спонтан бактериал перитонит билан асоратланишнинг профилактикаси ва давосини такомиллаштириш // *Oriental Renaissance: Innovative, educational, natural and social sciences.* 2023; P.947-953.
13. Sanokulova Sitora Avazovna. (2023). Acute Kidney Injury in Critically Ill Cirrhotic Patients with Spontaneous Bacterial Peritonitis. // *Texas Journal of Medical Science,* 2023;25:65-70. Retrieved from <https://zienjournals.com/index.php/tjms/article/view/4502>
14. Oblokulov A.R., M.I.Mukhammadieva. (2022). Clinical and biochemical characteristics of liver cirrhosis patients of viral etiology with spontaneous bacterial peritonitis // *Academia Globe: Inderscience Research.* 2022; P.210-216.
15. Облокулов А., Мухаммадиева М. (2022). Клинико-лабораторная характеристика спонтанного бактериального перитонита при циррозе печени вирусной этиологии. // *Журнал вестник врача,* 2022;1(3):66-69. извлечено от https://inlibrary.uz/index.php/doctors_herald/article/view/2016

Entered 20.08.2024