




New Day in Medicine
Новый День в Медицине

NDM



TIBBIYOTDA YANGI KUN

Ilmiy referativ, marifiy-ma'naviy jurnal



AVICENNA-MED.UZ



ISSN 2181-712X.
EiSSN 2181-2187

2 (76) 2025

Сопредседатели редакционной коллегии:

**Ш. Ж. ТЕШАЕВ,
А. Ш. РЕВИШВИЛИ**

Ред. коллегия:

М.И. АБДУЛЛАЕВ
А.А. АБДУМАЖИДОВ
Р.Б. АБДУЛЛАЕВ
Л.М. АБДУЛЛАЕВА
А.Ш. АБДУМАЖИДОВ
М.А. АБДУЛЛАЕВА
Х.А. АБДУМАДЖИДОВ
Б.З. АБДУСАМАТОВ
М.М. АКБАРОВ
Х.А. АКИЛОВ
М.М. АЛИЕВ
С.Ж. АМИНОВ
Ш.Э. АМОНОВ
Ш.М. АХМЕДОВ
Ю.М. АХМЕДОВ
С.М. АХМЕДОВА
Т.А. АСКАРОВ
М.А. АРТИКОВА
Ж.Б. БЕКНАЗАРОВ (главный редактор)
Е.А. БЕРДИЕВ
Б.Т. БУЗРУКОВ
Р.К. ДАДАБАЕВА
М.Н. ДАМИНОВА
К.А. ДЕХКОНОВ
Э.С. ДЖУМАБАЕВ
А.А. ДЖАЛИЛОВ
Н.Н. ЗОЛотова
А.Ш. ИНОЯТОВ
С. ИНДАМИНОВ
А.И. ИСКАНДАРОВ
А.С. ИЛЬЯСОВ
Э.Э. КОБИЛОВ
А.М. МАННАНОВ
Д.М. МУСАЕВА
Т.С. МУСАЕВ
М.Р. МИРЗОЕВА
Ф.Г. НАЗИРОВ
Н.А. НУРАЛИЕВА
Ф.С. ОРИПОВ
Б.Т. РАХИМОВ
Х.А. РАСУЛОВ
Ш.И. РУЗИЕВ
С.А. РУЗИБОВЕВ
С.А.ГАФФОРОВ
С.Т. ШАТМАНОВ (Кыргызстан)
Ж.Б. САТТАРОВ
Б.Б. САФОВЕВ (отв. редактор)
И.А. САТИВАЛДИЕВА
Ш.Т. САЛИМОВ
Д.И. ТУКСАНОВА
М.М. ТАДЖИЕВ
А.Ж. ХАМРАЕВ
Д.А. ХАСАНОВА
А.М. ШАМСИЕВ
А.К. ШАДМАНОВ
Н.Ж. ЭРМАТОВ
Б.Б. ЕРГАШЕВ
Н.Ш. ЕРГАШЕВ
И.Р. ЮЛДАШЕВ
Д.Х. ЮЛДАШЕВА
А.С. ЮСУПОВ
Ш.Ш. ЯРИКУЛОВ
М.Ш. ХАКИМОВ
Д.О. ИВАНОВ (Россия)
К.А. ЕГЕЗАРЯН (Россия)
DONG JINCHENG (Китай)
КУЗАКОВ В.Е. (Россия)
Я. МЕЙЕРНИК (Словакия)
В.А. МИТИШ (Россия)
В.И. ПРИМАКОВ (Беларусь)
О.В. ПЕШИКОВ (Россия)
А.А. ПОТАПОВ (Россия)
А.А. ТЕПЛОВ (Россия)
Т.Ш. ШАРМАНОВ (Казахстан)
А.А. ЩЕГОЛОВ (Россия)
С.Н. ГУСЕЙНОВА (Азербайджан)
Prof. Dr. KURBANHAN MUSLUMOV (Azerbaijan)
Prof. Dr. DENIZ UYAK (Germany)

**ТИББИЁТДА ЯНГИ КУН
НОВЫЙ ДЕНЬ В МЕДИЦИНЕ
NEW DAY IN MEDICINE**

*Илмий-рефератив, маънавий-маърифий журнал
Научно-реферативный,
духовно-просветительский журнал*

УЧРЕДИТЕЛИ:

**БУХАРСКИЙ ГОСУДАРСТВЕННЫЙ
МЕДИЦИНСКИЙ ИНСТИТУТ
ООО «ТИББИЁТДА ЯНГИ КУН»**

Национальный медицинский
исследовательский центр хирургии имени
А.В. Вишневского является генеральным
научно-практическим
консультантом редакции

Журнал был включен в список журнальных
изданий, рецензируемых Высшей
Аттестационной Комиссией
Республики Узбекистан
(Протокол № 201/03 от 30.12.2013 г.)

РЕДАКЦИОННЫЙ СОВЕТ:

М.М. АБДУРАХМАНОВ (Бухара)
Г.Ж. ЖАРЫЛКАСЫНОВА (Бухара)
А.Ш. ИНОЯТОВ (Ташкент)
Г.А. ИХТИЁРОВА (Бухара)
Ш.И. КАРИМОВ (Ташкент)
У.К. КАЮМОВ (Тошкент)
Ш.И. НАВРУЗОВА (Бухара)
А.А. НОСИРОВ (Ташкент)
А.Р. ОБЛОКУЛОВ (Бухара)
Б.Т. ОДИЛОВА (Ташкент)
Ш.Т. УРАКОВ (Бухара)

2 (76)

2025

февраль

www.bsmi.uz

https://newdaymedicine.com E:

ndmuz@mail.ru

Тел: +99890 8061882

LOCAL IMMUNE STATUS IN THE INFLAMMATORY PROCESS IN THE UPPER RESPIRATORY TRACT IN PATIENTS INFECTED WITH COVID-19

Narzullaev N.U. <https://orcid.org/0000-0002-9130-5568>

Bukhara State Medical Institute named after Abu Ali ibn Sina, Uzbekistan, Bukhara, st. A. Navoi. 1 Tel: +998 (65) 223-00-50 e-mail: info@bsmi.uz

✓ Resume

In the period from 2020-2021, we examined COVID-19 infected 53 patients who applied to the infectious diseases department of the Bukhara Regional Infectious Diseases Medical Center. Of these, 32 people suffered from acute rhinosinusitis and 21 people-acute purulent tonsillitis. All patients underwent a standard examination: clarification of complaints and anamnestic data. Examination of the upper respiratory tract organs was carried out according to the generally accepted scheme.

We also studied the indicators of local immunity of the mucous membranes of the upper respiratory tract using immunohistochemical studies with antibodies to T-helper lymphocytes (CD4 +), suppressor T-lymphocytes (CD8 +) and B-lymphocytes (CD20) in patients of different groups.

Indicators of local immunity of the mucous membrane of the nasal cavity and pharynx in the first two weeks of the acute process correspond to an adequate response of local immunity to an infectious agent; the third week is characterized by an imbalance of local immune protection, which indicates a high activity of the inflammatory process, which is a milestone in the course of the disease.

Key words: acute rhinosinusitis, acute tonsillitis, T-lymphocytes, B-lymphocytes.

МЕСТНАЯ ИММУННАЯ СТАТУС ПРИ ВОСПАЛИТЕЛЬНОМ ПРОЦЕССЕ В ВЕРХНЕМ ДЫХАТЕЛЬНОМ ПУТЕЙ У БОЛЬНЫХ ЗАРАЖЕННЫХ COVID-19

Нарзуллаев Н.У. <https://orcid.org/0000-0002-9130-5568>

Бухарский государственный медицинский институт имени Абу Али ибн Сины, Узбекистан, г. Бухара, ул. А. Навои. 1 Тел: +998 (65) 223-00-50 e-mail: info@bsmi.uz

✓ Резюме

В период с 2020-2021 годов мы обследовали заражен COVID-19 53 пациента, обратившихся в инфекционном -отделение Бухарский областной инфекционной медицинский центр.

Из них 32 человека страдали острым риносинуситом и 21 человек-острым гнойным тонзиллитом. Все пациенты прошли стандартное обследование: уточнение жалоб и анамнестических данных. Обследование органов верхних дыхательных путей проводилось по общепринятой схеме.

Мы также изучили показатели местного иммунитета слизистых оболочек верхних дыхательных путей с помощью иммуногистохимических исследований с антителами к Т-хелперным лимфоцитам (CD4 +), супрессорным Т-лимфоцитам (CD8 +) и В-лимфоцитам (CD20) у пациентов разных групп.

Показатели местного иммунитета слизистой оболочки полости носа и глотки в первые две недели острого процесса соответствуют адекватной реакции местного иммунитета на инфекционный агент; третья неделя характеризуется дисбалансом местной иммунной защиты, что свидетельствует о высокой активности воспалительного процесса, являющегося этапным этапом в течении заболевания.

Ключевые слова: острый риносинусит, острый тонзиллит, Т-лимфоциты, В-лимфоциты.

COVID-19 БИЛАН ЗАРАРЛАНГАН БЕМОРЛАРДА ЮҚОРИ НАФАС ЙЎЛЛАРИ КАСАЛЛИКЛАРИ МАВЖУД БЕМОРЛАРДА МАҲАЛЛИЙ ИММУНИТЕТНИНГ ҲОЛАТИ

Нарзуллаев Н.У. <https://orcid.org/0000-0002-9130-5568>

Абу али ибн Сино номидаги Бухоро давлат тиббиёт институти Ўзбекистон, Бухоро ш.,
А.Навоий кўчаси. 1 Тел: +998 (65) 223-00-50 e-mail: info@bsmi.uz

✓ Резюме

Тегишириш мақсадида 2020-2021-йиллар оралигида COVID-19 инфекцияси билан касалланган ва Бухоро вилоят юқумли касалликлар тиббиёт марказининг бўлимларига мурожаат қилган 53 нафар бемор олинди. Улардан 32 нафари ўткир риносинусит ва 21 нафари ўткир йирингли тонзиллит билан касалланган. Барча беморлар стандарт тегиширувдан ўтказилди: беморларнинг шикоятлари ва анамнестик маълумотлари йиғилди. Юқори нафас йўллари аъзоларини тегишириш умумий қабул қилинган схема бўйича амалга оширилди. Шунингдек, турли гуруҳдаги беморларда Т-хелпер лимфоцитлар (CD4+), супрессор Т-лимфоцитлар (CD8 +) ва В-лимфоцитлар (CD20) учун антителалар билан иммуногистокимёвий тадқиқотлар ёрдамида ўрта қулоқ шиллиқ пардаларининг маҳаллий иммунитет кўрсаткичларини ўргандик. Ўткир риносинуситнинг дастлабки икки ҳафтасида унинг маҳаллий иммунитет кўрсаткичлари инфекцион агентга маҳаллий иммунитетнинг адекват жавобига мос келади; учинчи ҳафтада маҳаллий иммун ҳимоя номуносивблиги билан характерланади, бу яллиғланиш жараёнининг юқори фаоллигини кўрсатади.

Калит сўзлар: ўткир риносинусит, ўткир тонзиллит, Т-лимфоцитлар, В-лимфоцитлар.

Relevance

The body's defenses are determined by systemic and local factors. Local protection is provided by the integrity of the mucous membrane of the VDP (upper respiratory tract) of the oral cavity, the microbicidal properties of saliva and lymphoid tissue. The integrity of the VDP mucosa is the best physiological barrier to infection. The protective factors of a tear are determined not only by its mechanical properties, but also depend on the biological substances dissolved in it that can cause the lysis of foreign agents.

Such substances include lysozyme, immunoglobulin A, and a large number of interleukins with bactericidal and antiviral effects. A powerful factor of local protection is the secretory IgA contained in saliva [1,2,5,8,9]. Nonspecific immunity of the VDP mucosa is provided by cellular elements — neutrophils and macrophages, as well as secretory elements — various chemotaxis factors for inflammatory agents (interleukins, leukotrienes, etc.) Specific immunity is provided by lymphoid tissue contained in a significant amount around the oral cavity — in the form of diffuse infiltration or in the form of nodular clusters devoid of a closed connective tissue case. The cells providing specific immunity are T-lymphocytes and plasmocytes [3,4].

In most cases, non-specific and specific immune protective factors neutralize bacteria and viruses. However, microorganisms often overcome these barriers, penetrate into the internal environment of the body and cause disease. This is facilitated by the weakening of local immunity, in particular against the background of COVID-19 [2,6,7,10].

In this regard, **the purpose** of this research work was to study the local immunity of the VDP mucosa in patients with different duration of the inflammatory process in the VDP.

Materials and methods

The methodology of the work is based on a systematic and comprehensive data analysis.

In the period from 2020-2021, we examined COVID-19 infected 53 patients who applied to the infectious diseases department of the Bukhara Regional Infectious Diseases Medical Center. Of these, 32 people suffered from acute rhinosinusitis and 21 people-acute purulent tonsillitis.

All patients underwent a standard examination: clarification of complaints and anamnestic data. Examination of the upper respiratory tract organs was carried out according to the generally accepted scheme. We also studied the indicators of local immunity of the mucous membranes of the upper respiratory tract using immunohistochemical studies with antibodies to T-helper lymphocytes (CD4+), suppressor T-lymphocytes (CD8+) and B-lymphocytes (CD20) in patients of different groups.

Indicators of local immunity of the mucous membrane of the nasal cavity and pharynx in the first two weeks of the acute process correspond to an adequate response of local immunity to an infectious agent; the third week is characterized by an imbalance of local immune protection, which indicates a high activity of the inflammatory process, which is a milestone in the course of the disease. Among the examined patients there were 19 (35.8%) men and 34 (64.1%) women aged 25 to 45 years. The duration of the disease ranged from 2 days to 2 years or more.

Group 1 (5 patients 9.4%) – 1st week of the disease; group 2 (6 patients; 11.3%) - 2nd week of the disease; group 3 (6 patients; 11.3%) – 3rd week of the disease; group 4 (4 patients; 7.5%) – the 4th week of the disease; group 5 (32 children; 60.4%) - patients with a disease duration of 6 months or more. All patients underwent a standard examination: clarification of complaints and anamnestic data. The inspection of the organs of the VDP was carried out according to the generally accepted scheme. Statistical data analysis was performed using Microsoft Excel 2010 and STATISTICA 10.0.

We also studied the indicators of local immunity of the mucous membranes of the VDP using immunohistochemical studies with antibodies to helper T-lymphocytes (CD4+), suppressor T-lymphocytes (CD8+) and B-lymphocytes (CD20) in patients of different groups. Analyzing the data obtained, it can be stated that in the 1st group of patients (1st week of the disease), there was a tendency for a slight increase in CD8 + T-suppressor lymphocytes (17.0 ± 4.9) and B-lymphocytes (CD20) (2.6 ± 1.9), with simultaneously unchanged CD4 + T-lymphocytes-helper cells (32.4 ± 12.9), which is associated with an adequate response of local immunity to a foreign agent.

The ratio of CD4+ T helper lymphocytes to CD8+ T suppressor lymphocytes in patients of this group reflects the normal state of local immune protection. In the 2nd group of children (2nd week of the disease) in comparison with the 1st group, the indicators of CD4+ T-helper lymphocytes (32.4 ± 12.9 and 31.7 ± 8.1) remain at the same level as a week ago, while a significant increase in the indicators of CD8 + T-suppressor lymphocytes (17.0 ± 4.9 and 31.4 ± 9.2 , $p < 0.05$). B-lymphocytes (CD20) (2.6 ± 1.9 and 12.4 ± 7.3 , $p < 0.05$) tend to moderate growth compared to the previous period of the disease. During this period, the ratio of CD4 + T-helper lymphocytes to CD8 + T-suppressor lymphocytes tends to 1, which indicates a severe course of the inflammatory process and a high activity of the immune system aimed at combating a foreign agent.

Changes in the local immunity of the mucous membrane of the middle ear in the third group of children (3rd week of the disease) in comparison with the 2nd group are characterized by the fact that the indicators responsible for the development of acute inflammation, namely the indicators of CD4 + T-helper lymphocytes (31.7 ± 8.1 and 18.9 ± 9.2 ; $p < 0.05$) and CD8+ T-suppressor lymphocytes (31.4 ± 9.2 and 10.3 ± 6.5 ; $p < 0.05$) decrease, and the index of B-lymphocytes (CD20) (12.4 ± 7.3 and 11.7 ± 14.6) remains approximately at the same level as a week ago. When analyzing the ratio of CD4+ T-helper lymphocytes to CD8+T-suppressor lymphocytes, a sharp increase was revealed, almost corresponding to the level of this indicator at the 1st week of the disease, i.e. the normal state of local immune protection.

Thus, at this time there is a clear imbalance in the local immune defense, indicating, most likely, a great activity of the inflammatory process and being a turning point throughout the disease, leading in the future, in the absence of adequate, timely treatment, in the conditions of bacterial superinfection, to the transformation of the acute inflammatory process into a persistent, irreversible chronic, the inflammatory process.

In the 4th group of children (4th week of the disease), in comparison with the 3rd group, an interesting trend of changes in all indicators responsible for local immunity is revealed. Indicators of both T-lymphocytes (CD4+ T-helper lymphocytes (18.9 ± 9.2 and 151.3 ± 14.9 , $p < 0.05$), CD8+ T-suppressor lymphocytes (10.3 ± 6.5 and 107.1 ± 27.1 , $p < 0.05$), and B-lymphocytes (CD20) (11.7 ± 14.6 and 126.1 ± 42.3 ; $p < 0.05$) increase sharply at this time.

At the same time, the ratio of CD4+ T-helper lymphocytes to CD8+ T-suppressor lymphocytes is again lower than the values indicating the normal state of local immune protection, which indicates a severe course of the inflammatory process on the one hand, and on the other hand, the available

reserve in local immune protection. In children of the 5th group with a disease duration of 6 months or more in comparison with the 4th group, signs of suppression of the immune response were revealed, as indicated by an increase in B-lymphocytes (CD20) (126.0 ± 42.3 and 214.4 ± 114.4 , $p < 0.05$).

The existing excess of T-lymphocytes (CD4+ T-helper lymphocytes, CD8+ T-suppressor lymphocytes) and B-lymphocytes (CD20) compared with the onset of the disease indicates that, despite the prescription of the process and the presence of subjective signs of remission, the inflammatory process is not completed and, apparently, the disease acquires the features of an autoimmune pathology.

Confirmation of the above is the approximation of the ratio of CD4+ T-helper lymphocytes to CD8+ T-suppressor lymphocytes to an indicator corresponding to hyperactivity.

Conclusion

Thus, the indicators of local immunity of the mucous membrane of the upper respiratory tract in the first two weeks of ORS and OT correspond to an adequate response of local immunity to an infectious agent; the third week is characterized by an imbalance in local immune protection (indicators of CD4 T-lymphocytes-helper and CD8 T-lymphocytes -suppressors decrease by 40.4% and 67.2%, respectively, and the indicator of B-lymphocytes (CD20) remains approximately at the same level), which indicates a large activity of the inflammatory process, which is a milestone stage during the disease.

LIST OF REFERENCES:

1. Aripova T.U., Ismailova A.A. COVID-19: possibilities of immunotherapy :review //Journal of Theoretical and Clinical Medicine: scientific and practical medical journal /Academy of Sciences of the Republic of Uzbekistan. - Tashkent : Institute of Immunology of the Academy of Sciences of the Republic of Uzbekistan. 2020;3:8-12.
2. Assanovich M.A. Escitalopram in pharmacotherapy of mental disorders in patients with COVID-19 //Medical news. 2021;1(316):59-62.
3. Bazdyrev E.D. Coronavirus infection - an urgent problem of the XXI century //Complex problems of cardiovascular diseases. 2020;9(2):6-16.
4. Bachilo E.V. Mental health of the population during the COVID-19 pandemic //Journal of Neurology and Psychiatry named after C.C. Korsakov. 2020;120(10):130-136.
5. Bekzhanova O.E., Kayumova V.R., Alimova S.H. Clinical and pathogenetic aspects of olfaction and taste disorders in COVID-19. Tashkent State Dental Institute 2021; 67-74 pp.
6. Belyakov N.A., Rassokhin V.V., Yastrebova E.B. Coronavirus infection COVID-19. the nature of the virus, pathogenesis, clinical manifestations. message 1 //HIV infection and immunosuppression. 2020;12(1):7-21.
7. Lantsov A.A., Khmel'nitskaya N.M., Endaltseva E.B. Local immunity and treatment of patients with chronic purulent otitis media //News of otorhinolaryngology and logopathology. 1999;1:3-7.
8. Narzulaev N.U., Khamidova N.K., Mirzayeva M.R. Clinical and immunological study of the effect of various types of therapy on the course of allergic rhinitis in children with hymenolepidosis //Annals of the Romanian Society of Cell Biology 2021; 1900-1908 pp.
9. Narzullaev N.U., Mirzayeva M.R. Immunological features of infectious diseases etiology of mononucleosis caused by Epstein-Barr virus in children //International Journal of Pharmaceutical Research 2021; 2667-2671 pp.
10. Narzullaev N.U., Rakhmatov A.A. Cytokine profile in children with acute inflammation of the middle ear on the background of chronic active hepatitis //New day medicine Tashkent 2021;2(34):15-17. <https://newdayworldmedicine.com/en/article/203>

Entered 20.01.2025