




New Day in Medicine  
Новый День в Медицине

NDM



# TIBBIYOTDA YANGI KUN

Ilmiy referativ, marifiy-ma'naviy jurnal



AVICENNA-MED.UZ



ISSN 2181-712X.  
EiSSN 2181-2187

2 (76) 2025

**Сопредседатели редакционной коллегии:**

**Ш. Ж. ТЕШАЕВ,  
А. Ш. РЕВИШВИЛИ**

Ред. коллегия:

М.И. АБДУЛЛАЕВ  
А.А. АБДУМАЖИДОВ  
Р.Б. АБДУЛЛАЕВ  
Л.М. АБДУЛЛАЕВА  
А.Ш. АБДУМАЖИДОВ  
М.А. АБДУЛЛАЕВА  
Х.А. АБДУМАДЖИДОВ  
Б.З. АБДУСАМАТОВ  
М.М. АКБАРОВ  
Х.А. АКИЛОВ  
М.М. АЛИЕВ  
С.Ж. АМИНОВ  
Ш.Э. АМОНОВ  
Ш.М. АХМЕДОВ  
Ю.М. АХМЕДОВ  
С.М. АХМЕДОВА  
Т.А. АСКАРОВ  
М.А. АРТИКОВА  
Ж.Б. БЕКНАЗАРОВ (главный редактор)  
Е.А. БЕРДИЕВ  
Б.Т. БУЗРУКОВ  
Р.К. ДАДАБАЕВА  
М.Н. ДАМИНОВА  
К.А. ДЕХКОНОВ  
Э.С. ДЖУМАБАЕВ  
А.А. ДЖАЛИЛОВ  
Н.Н. ЗОЛОТОВА  
А.Ш. ИНОЯТОВ  
С. ИНДАМИНОВ  
А.И. ИСКАНДАРОВ  
А.С. ИЛЬЯСОВ  
Э.Э. КОБИЛОВ  
А.М. МАННАНОВ  
Д.М. МУСАЕВА  
Т.С. МУСАЕВ  
М.Р. МИРЗОЕВА  
Ф.Г. НАЗИРОВ  
Н.А. НУРАЛИЕВА  
Ф.С. ОРИПОВ  
Б.Т. РАХИМОВ  
Х.А. РАСУЛОВ  
Ш.И. РУЗИЕВ  
С.А. РУЗИБОЕВ  
С.А. ГАФФОРОВ  
С.Т. ШАТМАНОВ (Кыргызстан)  
Ж.Б. САТТАРОВ  
Б.Б. САФОЕВ (отв. редактор)  
И.А. САТИВАЛДИЕВА  
Ш.Т. САЛИМОВ  
Д.И. ТУКСАНОВА  
М.М. ТАДЖИЕВ  
А.Ж. ХАМРАЕВ  
Д.А. ХАСАНОВА  
А.М. ШАМСИЕВ  
А.К. ШАДМАНОВ  
Н.Ж. ЭРМАТОВ  
Б.Б. ЕРГАШЕВ  
Н.Ш. ЕРГАШЕВ  
И.Р. ЮЛДАШЕВ  
Д.Х. ЮЛДАШЕВА  
А.С. ЮСУПОВ  
Ш.Ш. ЯРИКУЛОВ  
М.Ш. ХАКИМОВ  
Д.О. ИВАНОВ (Россия)  
К.А. ЕГЕЗАРЯН (Россия)  
DONG JINCHENG (Китай)  
КУЗАКОВ В.Е. (Россия)  
Я. МЕЙЕРНИК (Словакия)  
В.А. МИТИШ (Россия)  
В.И. ПРИМАКОВ (Беларусь)  
О.В. ПЕШИКОВ (Россия)  
А.А. ПОТАПОВ (Россия)  
А.А. ТЕПЛОВ (Россия)  
Т.Ш. ШАРМАНОВ (Казахстан)  
А.А. ЩЕГОЛОВ (Россия)  
С.Н. ГУСЕЙНОВА (Азарбайджан)  
Prof. Dr. KURBANHAN MUSLUMOV(Azerbaijan)  
Prof. Dr. DENIZ UYAK (Germany)

**ТИББИЁТДА ЯНГИ КУН  
НОВЫЙ ДЕНЬ В МЕДИЦИНЕ**

**NEW DAY IN MEDICINE**

*Илмий-рефератив, маънавий-маърифий журнал  
Научно-реферативный,  
духовно-просветительский журнал*

**УЧРЕДИТЕЛИ:**

**БУХАРСКИЙ ГОСУДАРСТВЕННЫЙ  
МЕДИЦИНСКИЙ ИНСТИТУТ  
ООО «ТИББИЁТДА ЯНГИ КУН»**

Национальный медицинский  
исследовательский центр хирургии имени  
А.В. Вишневского является генеральным  
научно-практическим  
консультантом редакции

Журнал был включен в список журнальных  
изданий, рецензируемых Высшей  
Аттестационной Комиссией  
Республики Узбекистан  
(Протокол № 201/03 от 30.12.2013 г.)

**РЕДАКЦИОННЫЙ СОВЕТ:**

М.М. АБДУРАХМАНОВ (Бухара)  
Г.Ж. ЖАРЫЛКАСЫНОВА (Бухара)  
А.Ш. ИНОЯТОВ (Ташкент)  
Г.А. ИХТИЁРОВА (Бухара)  
Ш.И. КАРИМОВ (Ташкент)  
У.К. КАЮМОВ (Тошкент)  
Ш.И. НАВРУЗОВА (Бухара)  
А.А. НОСИРОВ (Ташкент)  
А.Р. ОБЛОКУЛОВ (Бухара)  
Б.Т. ОДИЛОВА (Ташкент)  
Ш.Т. УРАКОВ (Бухара)

**2 (76)**

**2025**

*февраль*

[www.bsmi.uz](http://www.bsmi.uz)

<https://newdaymedicine.com> E:

ndmuz@mail.ru

Тел: +99890 8061882

Received: 20.01.2025, Accepted: 03.02.2025, Published: 10.02.2025

UDC 616

## THE IMPACT OF VARIOUS FACTORS ON THE FORMATION AND DEVELOPMENT OF THE CRANIOFACIAL AREA IN CHILDREN WITH HEARING LOSS

Mukhamedova Z.R. Email: [muxamedova.zarifa@bsmi.uz](mailto:muxamedova.zarifa@bsmi.uz)  
<https://orcid.org/0009-0003-4274-9394>

Bukhara State Medical Institute named after Abu Ali ibn Sina, Uzbekistan, Bukhara,  
st. A. Navoi. 1 Tel: +998 (65) 223-00-50 e-mail: [info@bsmi.uz](mailto:info@bsmi.uz)

✓ *Resume*

*Studied magazines, materials of scientific conferences, as well as other information sources for the collection of reliable information about the actions of the craniofacial region in children with hearing loss.*

**Keywords:** craniofacial region, hearing loss, early postnatal ontogenesis, various factors.

## ВЛИЯНИЕ РАЗЛИЧНЫХ ФАКТОРОВ НА ФОРМИРОВАНИЕ И РАЗВИТИЕ ЧЕРЕПНО-ЛИЦЕВОЙ ОБЛАСТИ У ДЕТЕЙ С ПОТЕРЕЙ СЛУХА

Мухамедова З.Р. Email: [muxamedova.zarifa@bsmi.uz](mailto:muxamedova.zarifa@bsmi.uz)  
<https://orcid.org/0009-0003-4274-9394>

Бухарский государственный медицинский институт имени Абу Али ибн Сины, Узбекистан,  
г. Бухара, ул. А. Навои. 1 Тел: +998 (65) 223-00-50 e-mail: [info@bsmi.uz](mailto:info@bsmi.uz)

✓ *Резюме*

*Изучены журналы, материалы научных конференций, а также другие источники информации для сбора достоверной информации о действиях черепно-лицевой области у детей с потерей слуха.*

**Ключевые слова:** черепно-лицевая область, потеря слуха, ранний постнатальный онтогенез, различные факторы.

## ESHITISH QOBILIYATINI YO'QOTGAN BOLALARDA BOSH-YUZ SOHASINING SHAKLLANISHI VA RIVOJLANISHIGA TURLI OMILLARNING TA'SIRI

Muxamedova Z.R. Email: [muxamedova.zarifa@bsmi.uz](mailto:muxamedova.zarifa@bsmi.uz)  
<https://orcid.org/0009-0003-4274-9394>

Abu Ali ibn Sino nomidagi Buxoro davlat tibbiyot instituti, O'zbekiston, Buxoro sh.  
A. Navoiy kochasi 1 Tel: +998 (65) 223-00-50 e-mail: [info@bsmi.uz](mailto:info@bsmi.uz)

✓ *Rezyume*

*Eshitish qobiliyatini yo'qotgan bolalarda bosh-yuz sohasining xatti-harakatlari to'g "risida ishonchli axborot to'plash uchun jurnallar, ilmiy anjumanlar materiallari, shuningdek, boshqa axborot manbalari o'rganildi.*

*Kalit so'zlar: bosh-yuz sohasi, eshitish qobiliyatini yo'qotish, erta postnatal ontogenet, turli omillar.*

**P**urpose of the study: to study the impact of various factors on the development of the craniofacial region in children with hearing loss.

**Materials and methods.** Information sources devoted to changes in the craniofacial region in children with hearing loss and materials related to the study of the impact of various factors were used.

## Results and discussion

Today, according to the World Health Organization (WHO), hearing loss is one of the leading factors that can worsen a person's lifestyle. 5.3% of the world's population is hearing impaired (WHO, 2012). One in 1,000 newborns is born with grade 3 or 4 sensorineural hearing loss. According to the Ministry of Emergency Situations, the number of children with severe hearing impairment and deafness who applied in Uzbekistan in 2014 was 520, this figure was 2000 due to the introduction of a modern diagnostic method - audio screening [1,3].

According to statistics (2021) from the United Nations (UN), more than 5% of the world's population, or 430 million people, need rehabilitation to address "disabling" hearing loss (432 million adults and 34 million children). It is estimated that by 2050 more than 700 million people, or one in ten, will have disabling hearing loss [2,4].

"Disabling" is defined as hearing loss in the better ear that exceeds 35 decibels (dB). Nearly 80% of these people live in low- and middle-income countries. Hearing loss is more common among older people: more than 25% of people over the age of 60 years suffer from this problem [5,6].

A person who is unable to hear as well as a person with normal hearing—a hearing threshold of 20 dB or less in both ears—suffers from hearing loss. Hearing loss can be mild, moderate, severe, or profound. It can develop in one or both ears and make it difficult to hear spoken language or loud sounds [6,10].

The concept of "hearing loss" applies to people with hearing loss ranging from mild to severe. Typically, hearing impaired people communicate using spoken language and can use hearing aids, cochlear implants and other assistive devices, as well as subtitles to improve hearing [7,8].

"Deaf" people in most cases suffer from profound hearing loss, that is, they hear very poorly or not hear at all. These people often use sign language to communicate.

It has been established that in children with deafness or hearing loss in more than 80% of cases, the disease is first diagnosed at the age of less than one year. In half of them, the disease is congenital or hereditary, that is, it appears as a result of the pathology of pregnancy, childbirth, or is transmitted genetically [3,4].

Describing the characteristic features of hearing loss in childhood, otolaryngologists distinguish several groups of patients. The basis is the time of onset of deafness. This is due to the fact that the cortex of the temporal lobes - the final section of the auditory analyzer - develops in a "sounding" environment. Even if parts of the brain are damaged, the child has a chance to keep hearing: the neurons surrounding the pathological focus form new connections and take on the functions of the dead. This is the phenomenon of neuroplasticity.

There are several options for hearing loss in babies: a deaf-mute child; hearing-impaired child; late deaf child.

A deaf-mute child had a persistent, bilateral hearing impairment that arose before the formation of speech. He does not understand the addressed speech and seeks to establish communication through facial expressions, gestures, manipulations with objects.

Late deaf children lose their hearing after some kind of illness or taking ototoxic drugs. Even if hearing loss occurs after the formation of speech, the lack of rehabilitation measures at preschool age leads to intellectual decline. Over time, these children become depleted in vocabulary, pronunciation becomes incorrect. With a decrease, hearing loss at a later age, the prognosis is more favorable, but pronunciation suffers to one degree or another: articulation, intonation are disturbed, errors in stress occur, speech becomes slurred [1,9].

A hearing-impaired child perceives sounds, but the degree of impairment varies. At the debut of the disease up to 1 year, speech development suffers: the cooing gradually disappears, the baby switches to sign language. Speech acquisition is almost impossible. Signs of a later hearing loss are: poor vocabulary, word distortion, violation of the grammatical structure of the sentence, "blurred", fuzzy speech.

Early diagnosis of hearing loss in young children will allow you to start rehabilitation measures on time, not to miss the optimal time for installing a cochlear implant, reconstructive surgery.

Hearing loss can be first diagnosed at any age. Refusal of medical care leads to a worsening of the condition. Adult patients may lose their jobs or gradually lose self-care skills. Hearing loss in children often leads to speech disorders and developmental delays, in particular in the craniofacial area [3,12].

The craniofacial complex includes the head, face, and oral cavity and is the most distinctive of all human body structures, giving individuals unique features. Structures of the craniofacial complex such as the mandible, palate, temporomandibular joint (TMJ) and dentition offer valuable paradigms for studying development, structure, and function [5,6].

Craniofacial deafness syndrome is characterized by characteristic facial features and profound hearing loss [3,4].

The distinctive facial features of people with craniofacial deafness syndrome are the result of various anomalies in the development of the skull and face. Deaf children often have underdeveloped or absent nasal bones, resulting in a small nose, thin nostrils, and a flattened midface with a flat bridge of the nose. People with this disorder usually also have widely spaced eyes (ocular hypertelorism), constricted eye openings (narrowed palpebral fissures), a small upper jaw (maxillary hypoplasia), and a small mouth with pursed lips. [11,13].

People with this condition also have profound hearing loss, which is caused by abnormalities in the inner ear (sensory deafness). Hearing loss in these people is present from birth [7,8].

### **Conclusion**

The pathology of even a part of the sensory system, in this case hearing, leads to a disruption in the processes of perception and assimilation of information, a slowdown in the formation of conditioned reflex connections necessary in the process of motor activity, including a number of motor manifestations that do not require direct participation of the auditory analyzer. Damage to the auditory analyzer primarily affects the work of the vestibular and kinesthetic analyzers, which entails a delay in the formation of the upright function, impaired muscle tone and the ability to maintain balance, underdevelopment of the ability to orientate in space, difficulties in differentiating motor sensations and performing complexly coordinated movements, in insufficient development of clear, smooth and measured movements [9,10].

### **List of used literature**

1. Tarasova N.V., Galonsky V.G., Eleseeva O.A., Orthodontic aspects of children and adolescents with sensory deprivation of hearing //Journal of Siberian Medical Sciences. 2013.
2. Khasanova D. Structural and functional features of the spleen of rats in the norm and with the introduction of a genetically modified product. //Society and innovations 2021;2(4):114-122.
3. Khasanova D.A. (2017). Modern tools for improving the efficiency of regional innovation structures. //In problems of effective use of the scientific potential of society 2017; 112-121 pp.
4. Akhrorovna K.D. Medical Field Morphological Features of Human and Mammalian Spleen in Postnatal Ontogeny. //Journal NX, 2021;7(1):252-256.
5. Bloch-Zupan A, Goodman JR. Otodental syndrome. //Orphanet J Rare Dis. 2006;1:5. Published 2006 Mar 21. doi:10.1186/1750-1172-1-5.
6. Brancati F, Sarkozy A, Dallapiccola B. KBG syndrome. //Orphanet J Rare Dis. 2006;1:50. Published 2006 Dec doi:10.1186/1750-1172-1-50.
7. Klingenberg C, Andersen WH. CHARGE-syndromet [The CHARGE syndrome]. Tidsskr Nor Laegeforen. 2008 Jun 12;128(12):1401-5. Norwegian. PMID: 18552902. 11. Purcell PL, Sie KC, Edwards TC, Doyle DL, Neidt K. Identification of oral clefts as a risk factor for hearing loss during newborn hearing screening. //J Early Hear Detect Interv. 2018;3(1):21-28.
8. Rachael B. Should infants with a syndrome or craniofacial anomaly be screened for a hearing loss at birth 2018;2(1):32-39.
9. Rahbar R, Robson CD et al., Craniofacial, temporal bone, and audiologic abnormalities in the spectrum of hemifacial microsomia. //Arch Otolaryngol Head Neck Surg. 2001 Mar;127(3):265-71. doi: 10.1001/archotol.127.3.265. PMID: 11255470.

**Entered 20.01.2025**

