



New Day in Medicine
Новый День в Медицине

NDM



TIBBIYOTDA YANGI KUN

Ilmiy referativ, marifiy-ma'naviy jurnal



AVICENNA-MED.UZ



ISSN 2181-712X.
EiSSN 2181-2187

5 (79) 2025

Сопредседатели редакционной коллегии:

**Ш. Ж. ТЕШАЕВ,
А. Ш. РЕВИШВИЛИ**

Ред. коллегия:

М.И. АБДУЛЛАЕВ
А.А. АБДУМАЖИДОВ
Р.Б. АБДУЛЛАЕВ
Л.М. АБДУЛЛАЕВА
А.Ш. АБДУМАЖИДОВ
М.А. АБДУЛЛАЕВА
Х.А. АБДУМАДЖИДОВ
Б.З. АБДУСАМАТОВ
М.М. АКБАРОВ
Х.А. АКИЛОВ
М.М. АЛИЕВ
С.Ж. АМИНОВ
Ш.Э. АМОНОВ
Ш.М. АХМЕДОВ
Ю.М. АХМЕДОВ
С.М. АХМЕДОВА
Т.А. АСКАРОВ
М.А. АРТИКОВА
Ж.Б. БЕКНАЗАРОВ (главный редактор)
Е.А. БЕРДИЕВ
Б.Т. БУЗРУКОВ
Р.К. ДАДАБАЕВА
М.Н. ДАМИНОВА
К.А. ДЕХКОНОВ
Э.С. ДЖУМАБАЕВ
А.А. ДЖАЛИЛОВ
Н.Н. ЗОЛотова
А.Ш. ИНОЯТОВ
С. ИНДАМИНОВ
А.И. ИСКАНДАРОВ
А.С. ИЛЪЯСОВ
Э.Э. КОБИЛОВ
А.М. МАННАНОВ
Д.М. МУСАЕВА
Т.С. МУСАЕВ
М.Р. МИРЗОЕВА
Ф.Г. НАЗИРОВ
Н.А. НУРАЛИЕВА
Ф.С. ОРИПОВ
Б.Т. РАХИМОВ
Х.А. РАСУЛОВ
Ш.И. РУЗИЕВ
С.А. РУЗИБОЕВ
С.А. ГАФФОРОВ
С.Т. ШАТМАНОВ (Кыргызстан)
Ж.Б. САТТАРОВ
Б.Б. САФОЕВ (отв. редактор)
И.А. САТИВАЛДИЕВА
Ш.Т. САЛИМОВ
Д.И. ТУКСАНОВА
М.М. ТАДЖИЕВ
А.Ж. ХАМРАЕВ
Д.А. ХАСАНОВА
Б.З. ХАМДАМОВ
А.М. ШАМСИЕВ
А.К. ШАДМАНОВ
Н.Ж. ЭРМАТОВ
Б.Б. ЕРГАШЕВ
Н.Ш. ЕРГАШЕВ
И.Р. ЮЛДАШЕВ
Д.Х. ЮЛДАШЕВА
А.С. ЮСУПОВ
Ш.Ш. ЯРИКУЛОВ
М.Ш. ХАКИМОВ
Д.О. ИВАНОВ (Россия)
К.А. ЕГЕЗАРЯН (Россия)
DONG JINCHENG (Китай)
КУЗАКОВ В.Е. (Россия)
Я. МЕЙЕРНИК (Словакия)
В.А. МИТИШ (Россия)
В.И. ПРИМАКОВ (Беларусь)
О.В. ПЕШИКОВ (Россия)
А.А. ПОТАПОВ (Россия)
А.А. ТЕПЛОВ (Россия)
Т.Ш. ШАРМАНОВ (Казахстан)
А.А. ЩЕГОЛОВ (Россия)
С.Н. ГУСЕЙНОВА (Азербайджан)
Prof. Dr. KURBANHAN MUSLUMOV (Azerbaijan)
Prof. Dr. DENIZ UYAK (Germany)

ТИББИЁТДА ЯНГИ КУН НОВЫЙ ДЕНЬ В МЕДИЦИНЕ NEW DAY IN MEDICINE

*Илмий-рефератив, маънавий-маърифий журнал
Научно-реферативный,
духовно-просветительский журнал*

УЧРЕДИТЕЛИ:

**БУХАРСКИЙ ГОСУДАРСТВЕННЫЙ
МЕДИЦИНСКИЙ ИНСТИТУТ
ООО «ТИББИЁТДА ЯНГИ КУН»**

Национальный медицинский
исследовательский центр хирургии имени
А.В. Вишневского является генеральным
научно-практическим
консультантом редакции

Журнал был включен в список журнальных
изданий, рецензируемых Высшей
Аттестационной Комиссией
Республики Узбекистан
(Протокол № 201/03 от 30.12.2013 г.)

РЕДАКЦИОННЫЙ СОВЕТ:

М.М. АБДУРАХМАНОВ (Бухара)
Г.Ж. ЖАРЫЛКАСЫНОВА (Бухара)
А.Ш. ИНОЯТОВ (Ташкент)
Г.А. ИХТИЁРОВА (Бухара)
Ш.И. КАРИМОВ (Ташкент)
У.К. КАЮМОВ (Тошкент)
Ш.И. НАВРУЗОВА (Бухара)
А.А. НОСИРОВ (Ташкент)
А.Р. ОБЛОКУЛОВ (Бухара)
Б.Т. ОДИЛОВА (Ташкент)
Ш.Т. УРАКОВ (Бухара)

5 (79)

2025

май

www.bsmi.uz

<https://newdaymedicine.com> E:

ndmuz@mail.ru

Тел: +99890 8061882

Received: 20.04.2025, Accepted: 06.05.2025, Published: 10.05.2025

UDC 616. 61: 615-276:577. 95. -092

DETERMINATION OF THE MORPHOMETRIC PARAMETERS OF THE EPIDIDYMIS OF WHITE OUTBREED RATS IN AGE DYNAMICS

Namozov Farrukh Jumaevich, <https://orcid.org/0009-0009-2434-2922>
e-mail: namozov.farruh@bsmi.uz

Bukhara State Medical Institute named after Abu Ali Ibn Sina, Uzbekistan, Bukhara Sh., A. Navoi street. Phone: +998 (65) 223-00-50 e-mail: info@bsmi.uz

✓ Resume

In this article, the normative morphometric parameters of the testicles in the age dynamics of white rats are presented. Epididymal weight and dimensions: Small testicular mass (0.21 g) and its dimensions (length - 0.74 cm, thickness - 0.49 cm) are observed in newborn rats. As the rats get older, the mass, length and thickness of the epididymis increase. The highest values are achieved in 12-month-old rats: weight - 0.76 g, length - 1.3 cm, thickness - 0.9 cm. Epididymal tissue volume and density: Epididymal tissue volume also increases with age, reaching the highest values in 12-month-old rats (0.64 cm³). The density of testicular tissue decreases with age, which may be related to tissue development and changes in its composition. Correlation between body weight and testicular weight: Both body weight and testicular weight increase with age of rats, which shows some parallelism in their growth.

Key words: testicular hypertrophy, purebred rats, morphometry, index.

ОПРЕДЕЛЕНИЕ МОРФОМЕТРИЧЕСКИХ ПАРАМЕТРОВ ПРИДАТКА ЯИЧЕК БЕЛЫХ БЕСПОРОДНЫХ КРЫС В ВОЗРАСТНОЙ ДИНАМИКЕ

Намозов Фаррух Джумаевич, <https://orcid.org/0009-0009-2434-2922>
e-mail: namozov.farruh@bsmi.uz

Бухарский государственный медицинский институт имени Абу Али ибн Сино, Узбекистан, Бухарский ш., ул. А. Навои. Тел.: +998 (65) 223-00-50 e-mail: info@bsmi.uz

✓ Резюме

В данной статье представлены нормативные морфометрические параметры яичек в возрастной динамике белых крыс. Масса и размеры придатка яичка: У новорожденных крыс наблюдается небольшая масса яичка (0,21 г) и его размеры (длина - 0,74 см, толщина - 0,49 см). По мере взросления крыс масса, длина и толщина придатка яичка увеличиваются. Наибольшие значения достигаются у 12-месячных крыс: вес - 0,76 г, длина - 1,3 см, толщина - 0,9 см. Объем и плотность придатка яичка: Объем придатка яичка также увеличивается с возрастом, достигая наибольших значений у 12-месячных крыс (0,64 см³). Плотность ткани яичка уменьшается с возрастом, что может быть связано с развитием ткани и изменением ее состава. Корреляция между массой тела и массой яичка: И масса тела, и масса яичка увеличиваются с возрастом крыс, что показывает некоторый параллелизм в их росте.

Ключевые слова: гипертрофия яичек, чистокровные крысы, морфометрия, индекс.

OQ ZOTSIZ KAMUSHLAR MOYAK ORTIG'INING MORFOMETRIK PARAMETRLARINI YOSH DINAMIKASIDA ANIQLASH

Namozov Farrux Jumaevich, <https://orcid.org/0009-0009-2434-2922>
e-mail: namozov.farruh@bsmi.uz

Abu Ali Ibn Sino nomidagi Buxoro davlat tibbiyot instituti, O'zbekiston, Buxoro Sh., A.Navoiy ko'chasi. Telefon: +998 (65) 223-00-50 e-mail: info@bsmi.uz

✓ Rezyume

Ushbu maqolada oq kalamushlarning yosh dinamikasidagi moyaklar me'yoriy morfometrik parametrlari keltirilgan. Epididimaning vazni va o'lchamlari: Yangi tug'ilgan kalamushlarda kichik moyak massasi (0,21 g) va uning o'lchamlari (uzunligi - 0,74 sm, qalinligi - 0,49 sm) kuzatiladi. Kalamushlar yoshi ulg'aygan sari epididimning massasi, uzunligi va qalinligi ortadi. Eng yuqori ko'rsatkichlarga 12 oylik kalamushlarda erishiladi: vazni - 0,76 g, uzunligi - 1,3 sm, qalinligi - 0,9 sm. Epididim to'qimalarining hajmi va zichligi: Epididimal to'qimalarning hajmi ham yoshga qarab ortadi va 12 oylik kalamushlarda (0,64 sm³) eng yuqori ko'rsatkichlarga etadi. Moyak to'qimalarining zichligi yosh bilan kamayadi, bu to'qimalarning rivojlanishi va uning tarkibidagi o'zgarishlar bilan bog'liq bo'lishi mumkin. Tana vazni va moyak og'irligi o'rtasidagi bog'liqlik: kalamushlarning yoshi bilan tana vazni ham, moyak vazni ham ortadi, bu ularning o'sishida bir oz parallellikni ko'rsatadi.

Kalit so'zlar: moyak gipertrofiyasi, naslli kalamushlar, morfometriya, indeks.

Relevance of the study

All over the world, special attention is being paid to research aimed at improving the early detection, treatment and prevention of diseases of the male reproductive system, including hypertrophy, caused by various physical factors. Currently, studies of the male population in different countries show an increase in spermatogenesis disorders [1-3], a decrease in the average age of men with spermogram abnormalities and various reproductive diseases [4-6]. The structure of the causes of male infertility is as follows: varicocele (14.8%), hypogonadism (10.1%), urogenital infections (9.3%), history of cryptorchidism (8.4%), previous oncological diseases (7.8 %), immunological factors (3.9%), erectile dysfunction and ejaculation (2.4%), systemic diseases (2.2%), vascular obstruction (2.2%), testicular tumors (1.2 %), presence of somatic diseases (7.7%). In about 30% of cases, the true cause of male infertility cannot be determined, in which case it is diagnosed as "idiopathic infertility" [2, 4-5]. Targeted measures are being implemented in our country to radically improve the healthcare system and improve the quality of medical services for the population. For this purpose, a number of tasks were defined, including "increasing the efficiency, quality and accessibility of medical care, implementing comprehensive measures aimed at supporting a healthy lifestyle and preventing diseases, including a healthy lifestyle the formation of a system of medical standardization, the introduction of high-tech diagnostic methods and treatment by creating effective models of patronage and clinical examination.

The purpose of the study: to study the morphometry of the testis in purebred rats .

Research materials

An experimental study was conducted on the material obtained from the testicles of 35 new-born, 3, 6, 9, 12 months old male white rats.

Young animals					H is the total number of animals
So birth	3 months old	6 months old	9 months old	12 months old	
7	7	7	7	7	35

Animals in the morning, hungry to the stomach, ether unconsciousness under one in the moment decapitation method slaughtered _ Belly space since opening then the testicles excesses separate received and their mass, length, width, volume and tissues density was studied. Testicle of excesses each one's mass electricity on the scales measured, length and width millimeter tape with was measured. Testicle of excesses according to the volume formula:

$V = 0.123 \times n \times c^2$, that is where: n, c - suitable, of the testicle length and thickness 0.123 - constant coefficient. Received testicle excesses Bowin in solution fixed. Raised to concentration have ethyl from alcohol from the past then they _ Hot paraffin inserted, then 6-7 μ m standard thickness has been testicle excesses pieces prepared, they are oriented sagittally or frontally. Samples to van Gieson according to

hematoxylin and eosin with painted _ Ready histological drugs NLCD-307B binoculars microscope under checked (Novel, China).

Research results In the process of studying the morphometry of the epididymis in non-white rats, the following data were obtained on newborn non-white rats: epididymal weight: 45.1 ± 1.46 g, body length: $8, 5 \pm 0.09$ cm, epididymis mass: 0.21 ± 0.011 g, epididymis length: 0.74 ± 0.016 cm, epididymis thickness: 0.49 ± 0.016 cm, epididymis thickness: volume: 0.13 ± 0.009 cm³, epididymis density: 1.6 g/cm³.

These results provide insight into the characteristics of testicular hypertrophy in newborn white outbred rats.

The data obtained from the study of the morphometry of the epididymis in newborn white outbred rats provide important information about the development and characteristics of this organ in this type of genotype.

It is interesting to note that the weight of the testicle is an important indicator that can be related to its functional activity and the general condition of the animal. Other parameters such as the length, thickness and volume of the epididymis also play an important role in evaluating their structure and possible effects on reproductive processes.




Figure 1. Morphometric dimensions of the testis of a nonwhite male rat (hematoxylin and eosin stain, medium magnification)




Figure 2. Morphometric dimensions of the testis of a nonwhite male rat (hematoxylin and eosin stain, medium magnification).

The analysis of the density of the epididymal tissue allows to understand the structural composition of this organ and can be useful in studying pathologies or the effect of various factors on the epididymal tissue. These results not only add important insights to the field of reproductive biology, but may also have practical implications for developing strategies to treat or prevent reproductive disorders in rats and possibly other mammals.

Morphometric parameters of epididymis of 3-month-old rats: epididymis body weight: 53 ± 0.65 g, body length: 9.1 ± 0.08 cm, epididymis mass: 0.32 ± 0.015 g, the length of the epididymis: 0.85 ± 0.018 cm, the thickness of the epididymis: 0.57 ± 0.014 cm, the volume of the epididymis: 0.20 ± 0.01 cm³, the tissue of the epididymis density: 1.64 g/cm³.

During the experiment with rats of different age groups, the following characteristics were revealed. In 6-month-old rats, the body weight of the testis is 69.5 ± 1.43 g, the body length is 10.7 ± 0.11 cm, the weight of the testis in them is 0.43 ± 0.015 g, The testis is the length of is calculated. 0.93 ± 0.020 cm, testicle excess thickness - 0.61 ± 0.015 cm, testicle excess volume - 0.25 ± 0.01 cm³, equal to g/cm³ (Figure 1). Testicle excess tissues density - 1.7 g/cm³ - organize did - 9 months old in rats testicle body weight of the rest reached 75.4 ± 1.35 g, body length reached 12.5 ± 0.39 cm. Testicle more weight 0.56 ± 0.020 g, testicle more length 0.98 ± 0.032 cm, testicle more thickness - 0.65 ± 0.011 cm, testicle more volume - 0.29 ± 0.016 cm³, testicle more tissues density g/cm³ - 1.8 (Figure 1). 12 months old in rats testicle the body weight of the rest is up to 101.4 ± 2.05 g increased, the body length reached 14.6 ± 0.34 cm. Testicle more weight 0.76 ± 0.025 g, testicle more length is equal to 1.3 ± 0.04 cm, testicle excess thickness - 0.9 ± 0.08 cm, testicle excess volume - 0.64 ± 0.02 cm³, testicle excess of tissues density was estimated as - 1.18 in g/cm³. Above morphometric check results given in Table 1 below.

Table 1

White without breed of rats to age circle normative morphometric indicators

Age (day)	Body mass (g)	Body length (cm)	Testicle more Weight (g)	testicle more length (cm)	testicle more thickness (cm)	testicle excess volume cm ³	Testicle excess of tissues density g/cm ³
new born	45.1 ± 1.46	8.5 ± 0.09	0.21 ± 0.011	0.74 ± 0.016	0.49 ± 0.016	0.13 ± 0.009	1.6
3 months old	53 ± 0.65	9.1 ± 0.08	0.32 ± 0.015	0.85 ± 0.018	0.57 ± 0.014	0.20 ± 0.01	1.64
6 months old	69.5 ± 1.43	10.7 ± 0.11	0.43 ± 0.015	0.93 ± 0.020	0.61 ± 0.015	0.25 ± 0.01	1.7
9 months old	75.4 ± 1.35	12.5 ± 0.39	0.56 ± 0.020	0.98 ± 0.032	0.65 ± 0.011	0.29 ± 0.016	1.8
12 months old	101.4 ± 2.05	14.6 ± 0.34	0.76 ± 0.025	1.3 ± 0.04	0.9 ± 0.08	0.64 ± 0.02	1.18

Young increase with body weight in rats and body length significant level increases. Young increase with Testicle excess of mass and of size increase is also correct is proportional. With that together, testicle excess al tissue density also increases, this young increase with of tissues structure and contained changes to show can

This results of rats development and physiology according to next studies for, as well as in mammals of tissues evolution and growth to understand for effect to do can

Of development different periods of rats physiological parameters comparative analysis to do interesting information determination enable will give. For example, age depending on body weight and testicle excess of size increase to adulthood reach processes and of rats in the body hormonal changes with depends to be can

With age testicle excess of tissues density also pay attention to the change focus it is necessary of density increase of tissues structure and their functional in activity changes to show can

Nutrition, light and temperature such as the environment of factors of rats physiological parameters taking into account the effect get need Various different development periods of rats physiological

parameters learning their biology and physiology about valuable information will give. Received data medicine, biology and pharmacology in the fields next studies for useful to be can

Summary

" White without breed in rats testicle more from the study of " morphometry ". the following conclusions release can:

1. Testicle more weight and dimensions:

- New born in rats testicle more small mass (0.21 g) and his dimensions (length - 0.74 cm, thickness - 0.49 cm) are observed.

- Rats aging with Testicle excess of mass, its length and thickness increases. Most high 12 months to indicators age in rats obtained: weight - 0.76 g, length - 1.3 cm, thickness - 0.9 cm.

2. Testicle excess tissues volume and density:

- Testicle more size and age looking increased, 12 months in rats the most high to indicators reaches (0.64 cm³).

- Testicle excess tissues density of age growth with decreases, this of tissues development and his of the composition change with depends to be can _

3. Body weight and testicle gain weight _ between dependency:

- Rats age with body weight and testicles excess the weight also increases, this their in growth some kind of parallelism shows.

This data reproductive biology and testicle more development and of activity medical aspects in the field next studies for useful to be can

LIST OF REFERENCES:

1. Epifanova MV, Epifanov AA, Artemenko SA A method of protection and restoration of spermatogenesis during surgical interventions on the spermatic cord, testicle, epididymis. - 2020. (in Russian).
2. Karimov Kh Ya. Anatomy - physiological features of the male reproductive system and the problem of male infertility. Monograph / Kh. Yes. Karimov, SA Ten, Sh. Zh . Teshaev . - T.: Uzbekistan national encyclopedia, 2011; 236 p. (in Russian).
3. Kashchenko SA., Zakharov AA Changes in the structure of the seminal vesicles of rats of the reproductive period under conditions of cyclophosphamide-induced immunosuppression. Yakut Medical Journal 2018;2:76-78. DOI: 10.25789/YMJ/2018. 62.22. (in Russ).
4. Kashchenko SA., Zakharov AA Organometric and morphometric changes in the testes of immature rats as a result of artificial immunosuppression // Zhurn. Honey. - biol. research. 2017;5(1):63-71. DOI: 10. 17238/issn2542-1298. 2017. 5. 1. 63. (in Russ).
5. Ghezzi M, De Toni L, Palego P, Menegazzo M, Faggian E, Berretta M, Fiorica F, De Rocco Ponce M, Foresta C, Garolla A. Increased risk of testis failure in testicular germ cell tumor survivors undergoing radiotherapy. Oncotarget. 2017 Dec 7;9(3):3060-3068.
6. Gong EJ, Shin IS, Son TG et al. Low-dose-rate radiation exposure leads to testicular damage with decreases in DNMT1 and HDAC1 in the murine testis. // J Radiat Res. 2014 Jan 1;55(1):54-60.

Entered 20.04.2025