



New Day in Medicine
Новый День в Медицине

NDM



TIBBIYOTDA YANGI KUN

Ilmiy referativ, marifiy-ma'naviy jurnal



AVICENNA-MED.UZ



ISSN 2181-712X.
EiSSN 2181-2187

5 (79) 2025

Сопредседатели редакционной коллегии:

**Ш. Ж. ТЕШАЕВ,
А. Ш. РЕВИШВИЛИ**

Ред. коллегия:

М.И. АБДУЛЛАЕВ
А.А. АБДУМАЖИДОВ
Р.Б. АБДУЛЛАЕВ
Л.М. АБДУЛЛАЕВА
А.Ш. АБДУМАЖИДОВ
М.А. АБДУЛЛАЕВА
Х.А. АБДУМАДЖИДОВ
Б.З. АБДУСАМАТОВ
М.М. АКБАРОВ
Х.А. АКИЛОВ
М.М. АЛИЕВ
С.Ж. АМИНОВ
Ш.Э. АМОНОВ
Ш.М. АХМЕДОВ
Ю.М. АХМЕДОВ
С.М. АХМЕДОВА
Т.А. АСКАРОВ
М.А. АРТИКОВА
Ж.Б. БЕКНАЗАРОВ (главный редактор)
Е.А. БЕРДИЕВ
Б.Т. БУЗРУКОВ
Р.К. ДАДАБАЕВА
М.Н. ДАМИНОВА
К.А. ДЕХКОНОВ
Э.С. ДЖУМАБАЕВ
А.А. ДЖАЛИЛОВ
Н.Н. ЗОЛотова
А.Ш. ИНОЯТОВ
С. ИНДАМИНОВ
А.И. ИСКАНДАРОВА
А.С. ИЛЬЯСОВ
Э.Э. КОБИЛОВ
А.М. МАННАНОВ
Д.М. МУСАЕВА
Т.С. МУСАЕВ
М.Р. МИРЗОЕВА
Ф.Г. НАЗИРОВ
Н.А. НУРАЛИЕВА
Ф.С. ОРИПОВ
Б.Т. РАХИМОВ
Х.А. РАСУЛОВ
Ш.И. РУЗИЕВ
С.А. РУЗИБОЕВ
С.А.ГАФФОРОВ
С.Т. ШАТМАНОВ (Кыргызстан)
Ж.Б. САТТАРОВ
Б.Б. САФОЕВ (отв. редактор)
И.А. САТИВАЛДИЕВА
Ш.Т. САЛИМОВ
Д.И. ТУКСАНОВА
М.М. ТАДЖИЕВ
А.Ж. ХАМРАЕВ
Д.А. ХАСАНОВА
Б.З. ХАМДАМОВ
А.М. ШАМСИЕВ
А.К. ШАДМАНОВ
Н.Ж. ЭРМАТОВ
Б.Б. ЕРГАШЕВ
Н.Ш. ЕРГАШЕВ
И.Р. ЮЛДАШЕВ
Д.Х. ЮЛДАШЕВА
А.С. ЮСУПОВ
Ш.Ш. ЯРИКУЛОВ
М.Ш. ХАКИМОВ
Д.О. ИВАНОВ (Россия)
К.А. ЕГЕЗАРЯН (Россия)
DONG JINCHENG (Китай)
КУЗАКОВ В.Е. (Россия)
Я. МЕЙЕРНИК (Словакия)
В.А. МИТИШ (Россия)
В.И. ПРИМАКОВ (Беларусь)
О.В. ПЕШИКОВ (Россия)
А.А. ПОТАПОВ (Россия)
А.А. ТЕПЛОВ (Россия)
Т.Ш. ШАРМАНОВ (Казахстан)
А.А. ЩЕГОЛОВ (Россия)
С.Н. ГУСЕЙНОВА (Азербайджан)
Prof. Dr. KURBANHAN MUSLUMOV (Azerbaijan)
Prof. Dr. DENIZ UYAK (Germany)

**ТИББИЁТДА ЯНГИ КУН
НОВЫЙ ДЕНЬ В МЕДИЦИНЕ
NEW DAY IN MEDICINE**

*Илмий-рефератив, маънавий-маърифий журнал
Научно-реферативный,
духовно-просветительский журнал*

УЧРЕДИТЕЛИ:

**БУХАРСКИЙ ГОСУДАРСТВЕННЫЙ
МЕДИЦИНСКИЙ ИНСТИТУТ
ООО «ТИББИЁТДА ЯНГИ КУН»**

Национальный медицинский
исследовательский центр хирургии имени
А.В. Вишневского является генеральным
научно-практическим
консультантом редакции

Журнал был включен в список журнальных
изданий, рецензируемых Высшей
Аттестационной Комиссией
Республики Узбекистан
(Протокол № 201/03 от 30.12.2013 г.)

РЕДАКЦИОННЫЙ СОВЕТ:

М.М. АБДУРАХМАНОВ (Бухара)
Г.Ж. ЖАРЫЛКАСЫНОВА (Бухара)
А.Ш. ИНОЯТОВ (Ташкент)
Г.А. ИХТИЁРОВА (Бухара)
Ш.И. КАРИМОВ (Ташкент)
У.К. КАЮМОВ (Тошкент)
Ш.И. НАВРУЗОВА (Бухара)
А.А. НОСИРОВ (Ташкент)
А.Р. ОБЛОКУЛОВ (Бухара)
Б.Т. ОДИЛОВА (Ташкент)
Ш.Т. УРАКОВ (Бухара)

5 (79)

2025

май

www.bsmi.uz

https://newdaymedicine.com E:

ndmuz@mail.ru

Тел: +99890 8061882

Received: 20.04.2025, Accepted: 06.05.2025, Published: 10.05.2025

UDC 616.34-008.11-022-053.2:577.21-07-08(575.1).

CURRENT CHALLENGES OF VIRAL DIARRHEA: DIAGNOSIS AND PREVENTION

Kholiev Bakhodir Oltibayevich

<https://orcid.org/0009-0005-9841-9145> e-mail: xoliyev.baxodir@bsmi.uz

Bukhara State Medical Institute named after Abu Ali ibn Sina, Uzbekistan, Bukhara, st. A. Navoi. 1
Tel: +998 (65) 223-00-50 e-mail: info@bsmi.uz

✓ Resume

Viral diarrhea remains a significant public health challenge worldwide, especially among children under five years of age. Despite advancements in diagnostic technologies and vaccination efforts, viral gastroenteritis caused by rotavirus, norovirus, adenovirus, and astrovirus continues to contribute to high morbidity and mortality rates, particularly in low- and middle-income countries. This review highlights the current epidemiological trends, diagnostic difficulties, and preventive strategies associated with viral diarrhea, emphasizing the need for integrated surveillance systems, improved sanitation, and global access to vaccination.

Keywords: *Viral diarrhea, rotavirus, norovirus, child health, epidemiology, vaccination, water sanitation.*

Relevance

Acute diarrheal diseases remain among the top causes of morbidity and mortality worldwide, particularly affecting children under five years of age [1]. Viral diarrhea, caused by a group of highly contagious enteric viruses, accounts for a substantial portion of these cases [2]. The most commonly implicated viruses include rotavirus, norovirus, adenovirus, and astrovirus [3]. These pathogens are transmitted primarily via the fecal-oral route, often through contaminated food, water, or direct person-to-person contact [4].

Despite the global rollout of rotavirus vaccines and improvements in hygiene awareness, viral diarrhea continues to cause significant health and economic burdens, especially in low- and middle-income countries. The persistence of these infections is closely linked to challenges such as inadequate sanitation, limited vaccine coverage, malnutrition, and weak healthcare infrastructure [5].

In recent years, the epidemiology of viral diarrhea has shifted, with a relative decline in rotavirus cases due to vaccination, and a rise in norovirus-associated outbreaks. Furthermore, the emergence of new viral strains, the lack of specific antiviral treatments, and diagnostic limitations have complicated the management and control of these infections [6].

This article aims to review the current challenges related to viral diarrhea, focusing on epidemiological trends, diagnostic limitations, and preventive strategies, while emphasizing the need for integrated public health interventions [7].

Aim of the Study: The aim of this study is to analyze the current challenges associated with viral diarrhea, with a particular focus on its epidemiology, diagnostic limitations, and preventive strategies. This research seeks to highlight the global burden of viral diarrheal diseases, identify gaps in current healthcare responses, and emphasize the importance of improved vaccination coverage, sanitation, and surveillance systems in controlling the spread of viral gastroenteritis. By synthesizing recent scientific findings and public health data, the study aims to support the development of more effective and sustainable interventions to reduce morbidity and mortality caused by viral diarrhea, particularly in vulnerable populations.

Materials and Methods

This study is a narrative review based on the analysis of recent scientific literature and global health reports related to viral diarrhea. Relevant data were collected from peer-reviewed journals, publications from the World Health Organization (WHO), Centers for Disease Control and Prevention (CDC), and other reputable databases such as PubMed, Scopus, and Google Scholar.

The literature search included articles published between 2015 and 2024, using keywords such as “viral diarrhea,” “rotavirus,” “norovirus,” “enteric viruses,” “childhood diarrhea,” “vaccine coverage,” “diagnostic challenges,” and “diarrhea prevention.” Studies focusing on epidemiological data, clinical features, diagnostic

methods, and public health interventions were included. Articles not available in English or lacking scientific validity were excluded.

Data were synthesized to identify current trends, challenges, and gaps in the diagnosis, treatment, and prevention of viral diarrhea. No primary data were collected from patients, and no ethical approval was required as this study is based solely on existing literature.

Results et discussion

The analysis of recent literature indicates that viral diarrhea remains a significant public health problem, particularly in low- and middle-income countries. Rotavirus continues to be the leading cause of severe diarrhea in children under five years of age, although a noticeable reduction in disease burden has been observed in countries with high vaccine coverage. In contrast, norovirus has emerged as a major cause of acute viral gastroenteritis in older children and adults, with frequent outbreaks reported in healthcare facilities, schools, and closed communities. Despite the availability of molecular diagnostic techniques such as PCR, many healthcare centers in resource-limited regions still rely on clinical diagnosis due to the high cost and lack of laboratory infrastructure. This results in underdiagnosis and misclassification of cases. Vaccination against rotavirus has been effective in reducing hospitalizations and deaths, but coverage remains insufficient in several countries, especially across sub-Saharan Africa and parts of Asia. Furthermore, no licensed vaccine currently exists for norovirus, which continues to contribute to a large number of unpreventable cases. Improvements in water supply, sanitation, and hygiene practices have shown a significant impact on reducing viral transmission, yet their implementation remains inconsistent. Overall, the reviewed data emphasize the urgent need for better access to diagnostics, increased vaccine coverage, and integrated public health interventions to reduce the burden of viral diarrhea.

Conclusion

Viral diarrhea remains a widespread and pressing global health issue, especially among children in low- and middle-income countries. Despite the introduction of rotavirus vaccines and improvements in hygiene and sanitation in many parts of the world, the disease continues to cause significant morbidity and mortality. Norovirus has become a growing concern due to its high infectivity and the absence of a licensed vaccine. Diagnostic limitations, uneven vaccine coverage, and poor access to clean water and sanitation facilities all contribute to the persistence of viral diarrheal diseases. Addressing these challenges requires a comprehensive public health approach that combines expanded immunization programs, improved diagnostic capabilities, and effective water, sanitation, and hygiene (WASH) interventions. Strengthening surveillance systems and increasing investment in vaccine development, particularly for norovirus, are essential steps toward reducing the global burden of viral diarrhea and improving health outcomes in vulnerable populations.

LIST OF REFERENCES:

1. Troeger, C., Khalil, I. A., Rao, P. C., Blacker, B. F., Clark, A., Reiner, R. C., ... Mokdad, A. H. (2018). Rotavirus vaccination and the global burden of rotavirus diarrhea among children younger than 5 years. *The Lancet Infectious Diseases*, 18(10), 1101-1110. [https://doi.org/10.1016/S1473-3099\(18\)30322-1](https://doi.org/10.1016/S1473-3099(18)30322-1)
2. Tate, J. E., Burton, A. H., Boschi-Pinto, C., Parashar, U. D., & WHO-coordinated Global Rotavirus Surveillance Network. (2016). Global, regional, and national estimates of rotavirus mortality in children under 5 years of age, 2000–2013. *Clinical Infectious Diseases*, 62(suppl_2), S96-S105. <https://doi.org/10.1093/cid/civ1013>
3. Ahmed, S. M., Hall, A. J., Robinson, A. E., Verhoef, L., Premkumar, P., Parashar, U. D., ... & Lopman, B. A. (2014). Global prevalence of norovirus in cases of gastroenteritis: A systematic review and meta-analysis. *The Lancet Infectious Diseases*, 14(8), 725–730. [https://doi.org/10.1016/S1473-3099\(14\)70767-4](https://doi.org/10.1016/S1473-3099(14)70767-4)
4. World Health Organization. (2023). Diarrhoeal disease. <https://www.who.int/news-room/fact-sheets/detail/diarrhoeal-disease>
5. Centers for Disease Control and Prevention (CDC). (2023). Global Diarrhea Burden. <https://www.cdc.gov/healthywater/global/diarrhea-burden.html>
6. Lopman, B. A., Steele, D., Kirkwood, C. D., & Parashar, U. D. (2016). The vast and varied global burden of norovirus: Prospects for prevention and control. *PLoS Medicine*, 13(4), e1001999. <https://doi.org/10.1371/journal.pmed.1001999>
7. Burnett, E., Parashar, U. D., & Tate, J. E. (2020). Global impact of rotavirus vaccination on diarrhea hospitalizations and deaths among children <5 years old: 2006–2019. *Journal of Infectious Diseases*, 222(10), 1731–1739. <https://doi.org/10.1093/infdis/jiaa081>

Entered 20.04.2025