



New Day in Medicine  
Новый День в Медицине

NDM



# TIBBIYOTDA YANGI KUN

Ilmiy referativ, marifiy-ma'naviy jurnal



AVICENNA-MED.UZ

ISSN 2181-712X.  
EiSSN 2181-2187

10 (84) 2025

**Сопредседатели редакционной  
коллегии:**

**Ш. Ж. ТЕШАЕВ,  
А. Ш. РЕВИШВИЛИ**

Ред. коллегия:  
М.И. АБДУЛЛАЕВ  
А.А. АБДУМАЖИДОВ  
Р.Б. АБДУЛЛАЕВ  
Л.М. АБДУЛЛАЕВА  
А.Ш. АБДУМАЖИДОВ  
М.А. АБДУЛЛАЕВА  
Х.А. АБДУМАДЖИДОВ  
Б.З. АБДУСАМАТОВ  
М.М. АКБАРОВ  
Х.А. АКИЛОВ  
М.М. АЛИЕВ  
С.Ж. АМИНОВ  
Ш.Э. АМОНОВ  
Ш.М. АХМЕДОВ  
Ю.М. АХМЕДОВ  
С.М. АХМЕДОВА  
Т.А. АСКАРОВ  
М.А. АРТИКОВА  
Ж.Б. БЕКНАЗАРОВ (главный редактор)  
Е.А. БЕРДИЕВ  
Б.Т. БУЗРУКОВ  
Р.К. ДАДАБАЕВА  
М.Н. ДАМИНОВА  
К.А. ДЕХКОНОВ  
Э.С. ДЖУМАБАЕВ  
А.А. ДЖАЛИЛОВ  
Н.Н. ЗОЛОТОВА  
А.Ш. ИНОЯТОВ  
С. ИНДАМИНОВ  
А.И. ИСКАНДАРОВ  
А.С. ИЛЬЯСОВ  
Э.Э. КОБИЛОВ  
А.М. МАННАНОВ  
Д.М. МУСАЕВА  
Т.С. МУСАЕВ  
М.Р. МИРЗОЕВА  
Ф.Г. НАЗИРОВ  
Н.А. НУРАЛИЕВА  
Ф.С. ОРИПОВ  
Б.Т. РАХИМОВ  
Х.А. РАСУЛОВ  
Ш.И. РУЗИЕВ  
С.А. РУЗИБОЕВ  
С.А. ГАФФОРОВ  
С.Т. ШАТМАНОВ (Кыргызстан)  
Ж.Б. САТТАРОВ  
Б.Б. САФОЕВ (отв. редактор)  
И.А. САТИВАЛДИЕВА  
Ш.Т. САЛИМОВ  
Д.И. ТУКСАНОВА  
М.М. ТАДЖИЕВ  
А.Ж. ХАМРАЕВ  
Б.Б. ХАСАНОВ  
Д.А. ХАСАНОВА  
Б.З. ХАМДАМОВ  
Э.Б. ХАККУЛОВ  
Г.С. ХОДЖИЕВА  
А.М. ШАМСИЕВ  
А.К. ШАДМАНОВ  
Н.Ж. ЭРМАТОВ  
Б.Б. ЕРГАШЕВ  
Н.Ш. ЕРГАШЕВ  
И.Р. ЮЛДАШЕВ  
Д.Х. ЮЛДАШЕВА  
А.С. ЮСУПОВ  
Ш.Ш. ЯРИКУЛОВ  
М.Ш. ХАКИМОВ  
Д.О. ИВАНОВ (Россия)  
К.А. ЕГЕЗАРЯН (Россия)  
DONG JINCHENG (Китай)  
КУЗАКОВ В.Е. (Россия)  
Я. МЕЙЕРНИК (Словакия)  
В.А. МИТИШ (Россия)  
В.И. ПРИМАКОВ (Беларусь)  
О.В. ПЕШИКОВ (Россия)  
А.А. ПОТАПОВ (Россия)  
А.А. ТЕПЛОВ (Россия)  
Т.Ш. ШАРМАНОВ (Казахстан)  
А.А. ІЦЕГОЛОВ (Россия)  
С.Н. ГУСЕЙНОВА (Азербайджан)  
Prof. Dr. KURBANHAN MUSLUMOV(Azerbaijan)  
Prof. Dr. DENIZ UYAK (Germany)

**ТИББИЁТДА ЯНГИ КУН  
НОВЫЙ ДЕНЬ В МЕДИЦИНЕ  
NEW DAY IN MEDICINE**

*Илмий-рефератив, маънавий-маърифий журнал  
Научно-реферативный,  
духовно-просветительский журнал*

**УЧРЕДИТЕЛИ:**

**БУХАРСКИЙ ГОСУДАРСТВЕННЫЙ  
МЕДИЦИНСКИЙ ИНСТИТУТ  
ООО «ТИББИЁТДА ЯНГИ КУН»**

Национальный медицинский  
исследовательский центр хирургии имени  
А.В. Вишневского является генеральным  
научно-практическим  
консультантом редакции

Журнал был включен в список журнальных  
изданий, рецензируемых Высшей  
Аттестационной Комиссией  
Республики Узбекистан  
(Протокол № 201/03 от 30.12.2013 г.)

**РЕДАКЦИОННЫЙ СОВЕТ:**

М.М. АБДУРАХМАНОВ (Бухара)  
Г.Ж. ЖАРЫЛКАСЫНОВА (Бухара)  
А.Ш. ИНОЯТОВ (Ташкент)  
Г.А. ИХТИЁРОВА (Бухара)  
Ш.И. КАРИМОВ (Ташкент)  
У.К. КАЮМОВ (Тошкент)  
Ш.И. НАВРУЗОВА (Бухара)  
А.А. НОСИРОВ (Ташкент)  
А.Р. ОБЛОКУЛОВ (Бухара)  
Б.Т. ОДИЛОВА (Ташкент)  
Ш.Т. УРАКОВ (Бухара)

**10 (84)**

www.bsmi.uz  
<https://newdaymedicine.com> E:  
ndmuz@mail.ru  
Тел: +99890 8061882

**2025  
октябрь**

Received: 20.09.2025, Accepted: 06.10.2025, Published: 10.10.2025

UDC 618.5-08:618.346-008.8

## ALGORITHM FOR THE STUDY OF CHILDBIRTH WITH PRENATAL RUPTURE OF AMNIOTIC FLUID, AGAINST THE BACKGROUND OF EROSION OF THE CERVIX

Khatamova M.T. <https://orcid.org/0000-0002-0279-0240>

Bukhara State Medical Institute named after Abu Ali ibn Sina, Uzbekistan, Bukhara, st. A. Navoi. 1  
Tel: +998 (65) 223-00-50 e-mail: [info@bsmi.uz](mailto:info@bsmi.uz)

✓ *Resume*

*Prenatal outflow of amniotic fluid combination with infectious diseases is a serious problem of modern obstetrics. The article presents modern views on the etiology and pathogenesis of premature and antenatal rupture of amniotic fluid. The pathological, genetic and microbiological aspects of this pathology are considered. Further solutions to the problems associated with premature and antenatal rupture of amniotic fluid are outlined. The causes of prenatal outflow of amniotic fluid, despite numerous studies, are not completely established, although the leading factor in this complication is considered to be infection.*

*Key words:* full-term pregnancy, induction of labor, prenatal outflow of amniotic fluid.

## АСПЕКТЫ ВЕДЕНИЯ РОДОВ ПРИ ДОРОДОВОМ ИЗЛИТИИ ОКОЛОПЛОДНЫХ ВОД, НА ФОНЕ ЭРОЗИИ ШЕЙКИ МАТКИ

Xатамова М.Т. <https://orcid.org/0000-0002-0279-0240>

Бухарский государственный медицинский институт имени Абу Али ибн Сины, Узбекистан, г. Бухара, ул. А. Навои. 1 Тел: +998 (65) 223-00-50 e-mail: [info@bsmi.uz](mailto:info@bsmi.uz)

✓ *Резюме*

*Дородовое и преждевременное излитие околоплодных вод, в сочетании с инфекционными заболеваниями и эрозий шейки матки является серьезной проблемой современного акушерства. В статье представлены современные взгляды на этиологию и патогенез преждевременного и дородового излития околоплодных вод. Рассмотрены патоморфологические, генетические и микробиологические аспекты данной патологии. Намечены дальнейшие пути решения проблем, связанных с преждевременным дородовым излитием околоплодных вод. Причины дородового и преждевременного излития околоплодных вод, несмотря на многочисленные исследования, окончательно не установлены, хотя ведущим фактором данного осложнения считают инфицирование.*

*Ключевые слова:* доношенный срок, индукция родов, дородовое излитие околоплодных вод, преждевременное излитие околоплодных вод.

## BACHADON BO'YNI EROZIYASI FONIDA AMNIOTIK SUYUQLIKNING MUDDATDAN OLDIN YORILISHIDA TUG'RUQ ASPEKTLARINI BOSHQARUVI

Xатамова М.Т. <https://orcid.org/0000-0002-0279-0240>

Abu Ali ibn Sino nomidagi Buxoro davlat tibbiyot instituti, O'zbekiston, Buxoro sh.  
A. Navoiy kochasi 1 Tel: +998 (65) 223-00-50 e-mail: [info@bsmi.uz](mailto:info@bsmi.uz)

✓ *Rezyume*

*Yuqumli kasalliklar va bachadon bo'yni eroziyasi bilan birligida amniotik suyuqlikning prenatal va erta yorilishi zamoniyligi akusherlikda jiddiy muammo hisoblanadi. Maqolada amniotik suyuqlikning erta va prenatal yorilishi etiologiyasi va patogenezi bo'yicha zamoniyligi qarashlar keltirilgan. Ushbu patologiyaning patomorfologik, genetik va mikrobiologik jihatlari ko'rib chiqiladi. Amniotik suyuqlikning prenatal va erta yorilishi bilan bog'liq muammolarni hal qilishning keyingi usullari ko'rsatilgan, ko'plab tadqiqotlarga qaramay, infektsiya bu asoratning asosiy omili hisoblanadi.*

*Kalit so'zlar:* to'liq muddat, tug'ruq induksiyasi, amniotik suyuqlikning prenatal yorilishi, amniotik suyuqlikning muddatidan oldin yorilishi.

### **Relevance**

One of the most common pregnancy complications is premature and prenatal discharge of amniotic fluid. There is no single point of view regarding the cause of premature rupture of the membranes. In the literature, the role and nature of changes in the structure of the membranes during prenatal and premature discharge of amniotic fluid are still being discussed. There is an assumption that the clinical options for premature discharge of water, as well as the features of the histological structure of the membranes, may determine differences in the degree of infection of the mother and child.

The study of the histological structure of the membranes showed that they are a metabolically active tissue and consist of amniotic epithelium, basement membrane, connective tissue, chorion and decidual membrane. Connective tissue is built from collagen types 1 and 3, which provide strength to the membranes. The basement membrane is located under the epithelium in the form of a narrow eosinophilic cell-free mass; the compact layer is represented by a homogeneous mass devoid of cells (indicating the strength of the amniotic membrane). The fibroblast layer is located in a dense network of collagen and reticular fibers and intercellular substance. The spongy layer of the amnion is connected through the connective tissue fibers and intercellular substance with a smooth chorion. Four layers are distinguished in the smooth chorion: cellular; reticular, containing fibroblasts, and a pseudobasal membrane formed by a trophoblast layer. The rupture of the membranes before the onset of labor is called the premature discharge of amniotic fluid (PIOV). Births complicated by premature discharge of amniotic fluid during full-term pregnancy are 15.1–19.6% and 5–35% during preterm delivery (up to 37 weeks of gestation) and have no tendency to decrease. Leading obstetrician-gynecologists note that this pathology contributes to the growth of complications in childbirth and in the postpartum period on the part of the mother, fetus and newborn. It should also be noted that PIOW tends to re-develop in subsequent births with a frequency of up to 20–32%. Factors leading to PIOW remain under discussion until now. Despite the constant attention of scientists to the PIOV problem, the etiology of this obstetric pathology remains not fully understood, there are no clear ideas about the possible mechanisms of rupture of the membranes. Ladhors L., Chernukha E. A., Savelyeva G.M., AriasF. consider that PIOV is a polyetiological pathology.

Prenatal amniotic fluid effusion (DIOD) is a serious problem of modern obstetrics. Amniotic fluid, or amniotic fluid, being a biologically active environment surrounding the fetus throughout pregnancy, perform a variety of functions, ensuring the normal functioning of the mother-placenta-fetus system. According to, labor against the background of prenatal rupture of the membranes is often accompanied by anomalies of labor, hypotonic and atonic bleeding, and high rates of trauma to the soft tissues of the birth canal.

Despite numerous studies, the causes of prenatal amniotic fluid outflow have not been conclusively established, although infection is considered the leading factor in this complication. Daneshmand et al., (2012) concluded that morpho-functional, physiological and biochemical changes in the genital tract during pregnancy lead to the fact that the vaginal microflora becomes more uniform, with a pronounced dominance of lactobacilli, which reduces the likelihood of contamination of the fetus conditionally pathogens during its passage through the birth canal. But childbirth leads to significant changes in the qualitative and quantitative composition of the vaginal microflora. Significantly increase the number of non-spore forming gram-negative strict anaerobes (mainly bacteroids), Escherichia, and levels of lactobacilli and bifidobacteria decrease. Violations of normal vaginal microflora contribute to the development of such an infectious complication as endometritis. One of the mechanisms for maintaining normal vaginal microflora is associated with the formation of lactobacilli during their metabolism of lactic acid and other organic acids that maintain a low pH of the vaginal environment. Acidification of various media during the growth of lactobacilli inhibits the proliferation of conditionally pathogenic microorganisms such as candida, peptostreptococci, bacteroids, gardnerella and other bacteria secreted from the vagina of women with dysbiotic disorders. Gram-negative obligate-anaerobic bacteria, some of their species, have pathogenic properties: they contain lipopolysaccharide in the cell wall, which is an inducer of IL-8, the main cytokine that triggers the inflammatory process. They are capable of producing succinic acid, which inhibits the migration of polymorphonuclear neutrophils and their phagocytic ability. Therefore, this increases the possibility of infection of the fetus and mother.

Prenatal outflow of amniotic fluid and prolongation of the anhydrous interval often leads to complications of the birth act (fast and rapid delivery, weakness and coordination of the contractile



activity of the uterus), which aggravates the condition of the fetus and in some cases requires surgical delivery. The frequency of prenatal rupture of the membranes varies widely: from 5 to 19.8% of cases of full-term pregnancy. Delivery in this case does not always end favorably for the fetus and mother. Childbirth and the postpartum period can have: the risk of purulent-septic complications in the mother, abnormalities of labor and intrauterine infection of the fetus. In addition, labor may be ineffective, which leads to an increase in the frequency of surgical interventions.

**Purpose of the study:** Studying the aspects of labor management in prenatal amniotic fluid outflows, at the same time studying the role of infection in DIA, to reduce obstetric and perinatal complications, and developing rational delivery management tactics in prenatal amniotic fluid discharge.

### **Materials and methods of examination:**

To solve the tasks, a comprehensive examination of 72 pregnant women was carried out, the birth of which was complicated by DIA in the periods of 37-40 weeks of gestation, received by the Bukhara regional perinatal center for the period of 2017. Using the anamnestic, clinical, laboratory and instrumental data, we studied the course of pregnancy, childbirth, the postpartum period, the condition of the fetus and the newborn. The readiness of the birth canal was assessed using the Bishop scale. According to the National Standard for the Management of Patients with DIA, after 18 hours of anhydrous interval, antibacterial therapy was carried out to prevent purulent-septic complications in puerperas. The birth canal in the childbirth DIOV examined after 24 hours in the absence of labor in order to resolve the issue of the advisability of induction of labor. The nature of labor was controlled on the basis of partograms. Conducted: monitoring hemodynamic parameters, maintaining an observation sheet, measuring to-body every 4 hours, laboratory monitoring of leukocytes 1 time per day, general urine analysis, analysis of vaginal discharge. Ultrasound of the uterus and the fetus, cervicometry, monitoring of the rhythm and heart rate of the fetus and the general condition of the woman in labor were performed. Given the high sensitivity to ampicillin of the bacteria of the vagina and cervix, this antibacterial drug was used according to the protocol. The condition of the fetus was evaluated according to ultrasound and cardiotocography (CTG), and the condition of the newborn at birth, according to the Apgar scale. Fetal monitoring in childbirth was performed using a Corometrics 170 apparatus.

### **Results and its discussion:**

The average age of women observed was 26.5 years. In all women, pregnancy proceeded against the background of extragenital diseases, and in most cases a combination of several of them. Anemia of mild to moderate severity (72.2%), thyroid disease (33.3%) and varicose veins (25%) predominated. Every third woman (32%) suffered an infectious disease during this pregnancy mainly in the form of ARI, exacerbation of chronic sinusitis, cystitis, pyelonephritis. In 16.7% of pregnant women, ARI episodes were repeated many times during pregnancy. Among the transferred gynecological diseases, colpitis of various etiologies, which accounted for 43%, are most often diagnosed. 72.2% of women had a history of inflammatory diseases of the genital tract. This was mainly manifested in the form of yeast, trichomonas and banal colpitis, endometritis and adnexitis. 19.4% of women were treated for cervicitis and cervical erosion. According to previous analyzes of vaginal smears, 43% of women had grade 3 and 4 purity of vaginal smears.

In all women with prenatal amniotic fluid, a vaginal examination was performed to assess the maturity of the cervix according to the Bishop scale. Assessment was carried out according to 5 criteria. It was revealed that 61.1% of the pregnant women examined had disclosure, length, consistency, position of the cervix and the condition of the pre-existing part of the fetus with scores of up to 5, which was assessed as "immature neck". And in 38.9% of women, the birth canal was assessed as a "mature neck".

Accordingly, the tactics of further conduct was chosen according to the OPC protocol. In pregnant women with an immature neck, the induction of labor by Glandin E 2, 3 mg, 1 tablet intravaginally after the informed consent of the pregnant woman and relatives is proposed. A discussion was held about possible complications of labor excitement. During induction, fetal heartbeats and uterine activity were monitored. The birth canal is reevaluated after 8 hours to clarify the need for continued induction. In pregnant women with a "mature" cervix, delivery was carried out with expectant tactics until regular

labor was played out, or a consultation of doctors resolved the issue of oxytocin stimulation. 58.3% of pregnant women were delivered through the natural birth canal. Newborns born to mothers with DIAIs were evaluated on an Apgar scale by an average of 6 points.

Thus, studies have shown that the majority of women in labor with prenatal rupture of the membranes, cervical preparedness was estimated at 5 points, which meant "unpreparedness" of the birth canal. Of this number, 58.3% of women in labor underwent childbirth after the informed consent of the woman in labor and her relatives. 22.2% of women in labor had relative or absolute contraindications to labor stimulation and birth control. The remaining 19.4% of women in labor refused delivery, who chose Caesarean section as a further tactic of delivery. A study of the postpartum period showed that 26.4% of women experienced complications such as a lohiometer and a hematometer, manifested in the form of uterine subinvolution, substantiated by clinical data and ultrasound studies. In 18.1% of women, secondary healing of soft birth canal wounds was observed. In 2.8% of women, the postpartum period was accompanied by signs of exacerbation of chronic inflammatory diseases of the genital tract.

### **Findings:**

1. A long anhydrous period is a factor in increasing infection, which leads to an increase in obstetric and perinatal pathology.
2. A long anhydrous period increases the incidence of newborns and puerperas in the postpartum period.
3. A long anhydrous period increases the seeding of the birth canal of the general and conditionally pathogenic flora and leads to an increase in the imbalance of the vaginal ecosystem.

These data dictate the need for the use of antibacterial drugs in women in labor with DIA. The appointment of antibiotic therapy after 18 hours of anhydrous interval prevents the development of purulent-inflammatory processes in the body of the woman in labor and the fetus, causes colonization resistance and is not a contraindication to surgical delivery by cesarean section.

### **LIST OF REFERENCES:**

1. Khatamova M.T. Agnostic criteria for prenatal rupture of amniotic fluid Научно-практический журнал Казахстан Акушерство, гинекология и перинатология 2021;1(85).
2. Khatamova M.T. Pharmacogenetic and morphological features of stomach cancer on the background of uterine endometriosis // Tibbiyotda yangi kun 2021;2(34/3):114
3. Khatamova M.T., Bishekova B.Z. Aspects of labor outcome in women suffering with chronic pyelonephritis // Tibbiyotda yangi kun 2021;2(35/1)489.
4. Fayzulloeva N.Sh., Khotamova M.T., Hemostasiological manifestations constant septic diseases // Tibbiyotda yangi kun 2021;2(34/3):114
5. Kalmataeva Zh.A., Khatamova M.T. Aspects of amenorhea of hypergonadotropic genesis // Tibbiyotda yangi kun 2021;2(34/3):476
6. Fayzulloeva N.Sh., Khotamova M.T. Aspects of kidney state in women of reproductive age for uncomplicated pregnancy // Tibbiyotda yangi kun 2021;2(34/3):486
7. Khatamova M.T., Faizulloeva N.Sh. Features of ovarian failure, gonadal dysgenesis // Tibbiyotda yangi kun 2021;2(34/3):501
8. Shukurlaeva Sh.J., Hotamova M.T., Fayzulloeva N.Sh. Application of nitroxaline in pregnant women and aspects of labor outcome in women suffering with chronic pyelonephritis // Tibbiyotda yangi kun 2021;2(34/1):107.
9. Khotamova M.T., Fayzulloeva N.Sh. Farmakogenelogical and morphological features of stomach cancer during Endoscopic examination // Tibbiyotda yangi kun 2021;2(34/1):134-138.

**Entered 20.09.2025**